

## REVIEW

# ***Salmonella enterica* serovar Enteritidis: a Mini-review of Contamination Routes and Limitations to Effective Control**

**Charles O. A. OMWANDHO<sup>1†</sup> and Takayuki KUBOTA<sup>2\*</sup>**

<sup>1</sup> Department of Biochemistry, University of Nairobi (P.O. Box 30197, Nairobi, Kenya)

<sup>2</sup> Department of Drug Formulation Development, National Institute of Animal Health, National Agriculture and Food Research Organization (Tsukuba, Ibaraki 305-0856, Japan)

### **Abstract**

Salmonellosis is associated with massive public health and economic losses globally. It is estimated to cost poultry farmers in the United States of America up to US\$114 million annually. Attempts to develop effective vaccines and eradicate *Salmonella enterica* serovar Enteritidis (*S. Enteritidis*) from hen houses are undermined by serious limitations. This article reviews documented contamination routes and limitations to the rapid development of vaccines. Host-parasite interactions and clinical pathology are discussed and methods for reducing *S. Enteritidis* infection and transmission suggested.

**Discipline:** Animal health

**Additional key words:** chicken, farm-house environment, vaccine development

## **Introduction**

Salmonellosis is associated with contaminated poultry products. It presents major public health and economic concerns globally. Previous reports indicate that up to 3.7 million cases of salmonellosis occur in the United States of America (USA) every year, with economic losses to poultry farmers estimated at US \$64 million to US \$114 million annually<sup>13</sup>. Attempts to develop effective vaccines and eradicate *Salmonella enterica* serovar Enteritidis from hen houses are undermined by serious limitations. To ensure safety of products and maximization of profits, there is a need to gain firm knowledge on improved breeding techniques to ensure protection against possible outbreaks of *Salmonella* food poisoning. This article reviews *S. Enteritidis* contamination routes and control, giving a fresh look at documented contamination routes and re-examines limitations to vaccine development. Host-parasite interactions and associated clinical pathology are discussed and methods for reducing *S. Enteritidis* infection and transmission suggested.

## ***Salmonella enterica* serovars associated with food poisoning**

*Salmonella enterica* is one of the most common causes of food poisoning in humans<sup>76</sup>. It has 6 subspecies and about 2,500 serotypes<sup>99</sup>, of which the two most frequently reported serovars are *Salmonella enterica* serovar Typhimurium (*S. Typhimurium*) and *S. Enteritidis*, a Gram-negative bacterium that negatively affects both human and animal health. Using the Colindale phage-typing scheme, 37 phages have been used to identify more than 210 phage types for *S. Typhimurium*<sup>2,3,15,29</sup>. On the other hand, 16 phages have been used to identify 65 phage types for *S. Enteritidis*<sup>97</sup>. Historically, *S. Typhimurium* was among the most common serovars isolated from poultry across many countries from the 1950s to the late 1970s, but was overtaken by *S. Enteritidis* as the most common serovar isolated from poultry in the mid 1980s to date<sup>75</sup>. In England and Wales, the percentage of *S. Enteritidis* isolated from poultry rose from 3.3% of all *Salmonella* serovars in 1985 to 47.8% in 1988 and 48.3% in 1989<sup>75</sup>. Of these, the *S. Enteritidis* phage type 4 (PT4) was the most frequently reported and accounted for 71% of the isolates in 1988<sup>62</sup>. A three month survey conducted in 1990

<sup>†</sup>Dr. Charles O. A. Omwandho was sponsored by The Japanese Society For Promotion of Science to take a ten month sabbatical (1-12-2006 to 30-09-2007) at the National Institute of Animal Health, Tsukuba.

\*Corresponding author: e-mail [tkubota@affrc.go.jp](mailto:tkubota@affrc.go.jp)

Received 30 September 2008; accepted 20 February 2009.

to estimate prevalence and distribution of *S. Enteritidis* in spent laying hens in commercial egg production flocks in the USA demonstrated the presence of *S. Enteritidis* in 3% of 23,431 pooled caecal samples collected from 406 layer houses<sup>26</sup>.

## **S. Enteritidis and the burden of salmonellosis in humans and poultry**

*Salmonella* Enteritidis has been associated with pandemics in Europe and elsewhere. Up to 3.7 million cases of salmonellosis were estimated to occur annually in the United States alone<sup>63</sup> with economic losses to poultry farmers estimated at US\$ 64 to 114 million due to salmonellosis in young chickens<sup>13</sup>. Persons infected with *S. Enteritidis* usually develop fever, abdominal cramps and diarrhea beginning 12 to 72 hours after consuming contaminated food. The illness often lasts 4 to 7 days and most persons recover without antibiotic treatment. However, in the elderly, infants and persons with impaired immune systems, diarrhea can be severe and the persons may be ill enough to require hospitalization. In such patients, infection may spread to other body organs and can cause death if prompt antibiotic treatment is not administered. The cost estimates per case of human salmonellosis range from approximately US \$40 to US \$4.6 million respectively for uncomplicated cases to those ending with hospitalization and death<sup>100</sup>. Epidemiological data in Hungary, United Kingdom, USA, and Germany have confirmed that the food most commonly associated with increased illness in humans was the egg<sup>72,85,90</sup> and that *S. Enteritidis* accounted for 85% of all cases of human salmonellosis in Europe<sup>30</sup>.

Cases of animal and human salmonellosis have been grossly underestimated due to lack of systematic methods of reporting<sup>27</sup>. Also, whether or not specific incidences are reported depends on intensity of surveillance<sup>34</sup>, submission of isolates for serotyping<sup>84</sup>, severity of illness and association with a recognized outbreak in human populations<sup>1,88,92</sup>. Thus, although outbreaks would easily attract media attention, 80% of all salmonellosis cases occur individually rather than as outbreaks<sup>100</sup>.

## **Host–parasite interactions and clinical symptoms in chickens**

The outcome of an interaction between *Salmonella* and its host is dependent upon multiple factors including the genetic background of the host<sup>51</sup>. Previous studies have revealed considerable differences between lines in the level of colonization of the gastrointestinal tract and response to vaccination<sup>57,59</sup>. Also, while salmonellosis in young chickens is characterized by severe clinical signs of systemic disease

associated with diarrhea, dehydration and high mortality rates<sup>93</sup>, adult chickens can produce *Salmonella*-contaminated eggs without evidence of discernible illness<sup>31</sup>. Poultry infected with other egg contaminating serotypes, such as *Salmonella enterica* serovar Pullorum and *Salmonella enterica* serovar Gallinarum, are associated with drastic weight loss, a sharp decrease in egg production, and increased mortality<sup>80,82</sup>. On the other hand, there are typically no clinical signs in birds infected with *S. Enteritidis*<sup>44</sup>.

## **Contamination of eggs with S. Enteritidis**

Previous reports indicate that *S. Enteritidis* is the only human pathogen that contaminates eggs routinely even though the chicken farm environment is a rich source of other *Salmonella* serotypes<sup>12,14,83</sup>. Also, while penetration of cracked eggshells by bacteria was seen as a frequent cause of human illness before the introduction of grading schemes in the 1970s, but it is now known that egg contamination by *S. Enteritidis* may occur by vertical transmission in the reproductive tract before deposition of the shell<sup>37,48,64</sup>. In one study, adult laying hens were inoculated orally with 10<sup>8</sup> colony-forming units (cfu) of *S. Enteritidis* and the bacterium was isolated 2 days post infection from spleen, liver, heart, gall bladder, intestinal tissues, and from various sections of the ovary and oviduct<sup>47,49</sup>. Additional reports indicate that eggs can be contaminated with faeces from hens excreting *Salmonella*<sup>11,39,79,86,89,101</sup>. In such cases, *Salmonella* in faeces are believed to penetrate egg shell pores as the egg cools and before the establishment of the proteinaceous cuticular barrier<sup>10,31,86,101</sup>. Fecal matter adherent to the shell may also contaminate eggs via cracks on egg shells or when eggs are broken open for preparation of food products<sup>11</sup>. Previous reports indicate that forming eggs are subjected to descending infections from colonized ovarian tissue, lateral infections from upper oviduct and ascending infections from colonized vaginal and cloacal tissues<sup>49</sup>. It has also been suggested that *S. Enteritidis* may actively gain access to the egg by penetrating marginally faulty shells that exclude most other bacteria<sup>5</sup>.

## **Horizontal infection of S. Enteritidis in poultry**

Horizontal transmissions occur during hatching of chicks, with spread of *S. Enteritidis* in aerosol contamination, litter, dust, and faecal as well as caecal droppings of littermates<sup>71</sup>. There is evidence that *S. Enteritidis* spreads rapidly from infected day old chicks to pen mates reared on litter<sup>78,81</sup>. Poultry may also get infected from feed<sup>43,66</sup>, water<sup>36</sup>, rodents<sup>4,17,40</sup>, or by contact with other poults or chicks. Infection by these routes in chicks reached 100% within seven days of contact<sup>81</sup>. Contamination through water

results from the fact that young chicks and poults often dip their beaks, walk in, and defaecate in drinking water before drinkers are raised from the floor of the barn. In one study, laying hens experimentally inoculated with  $10^5$  *S. Enteritidis* phage type 4 (PT4) spread the infection via drinking water to un-inoculated hens in only 1 to 5 days<sup>69</sup>. Further evidence shows that short-term exposure to environmental stresses, such as introduction of young chicks to the same rearing room, and moulting (removal of feed and water from laying hens for 2 days), is associated with an increase in *S. Enteritidis* shedding by laying hens<sup>69</sup>.

### Hen-house management and environmental considerations

The ubiquitous presence of *Salmonella* within the poultry shed environment provides opportunity for multiple modes of transmission<sup>18</sup>. However, management styles or the manner in which chickens are kept in poultry houses can influence the rate of infection. In one study, a lower prevalence of *S. Enteritidis* PT4 was reported among 176,000 caged laying hens compared to free-range hens (1.7% vs. 50%) and the prevalence in culled hens kept in dirt-floor houses ranged from 14% to 42%<sup>52</sup>. Exposure to airborne *S. Enteritidis* PT4 can lead to generalized infection, even at low exposure doses<sup>7</sup>. Also, *S. Enteritidis* PT4 can survive in aerosols up to 2 hours with a negligible reduction in numbers<sup>61</sup>. A report by Davies and Wray<sup>19</sup> indicated that *S. Enteritidis* can survive for at least one year in empty poultry sheds, where naturally infected flocks have been housed, but will decline rapidly in litter. Elsewhere, *S. Enteritidis* from human sewage effluent discharged upstream of a chicken farm infected chickens in a commercial farm in southern California<sup>19,20,53</sup>. Isolates from hens were of the same phage type (PT4) as those found in eggs, sewage effluent and mice trapped in hen houses, and from cats and skunks on the premises<sup>19,20,95</sup>. Another report revealed that *S. Enteritidis* and fifty other different *Salmonella* serovars were isolated from hen house environment samples (litter and water) in Canadian flocks<sup>74</sup>.

Infected domestic animals can become healthy carriers, latently infected or less frequently clinically ill, but may excrete *Salmonella* in their feces and form a large reservoir and source of environmental contamination for other animals and humans. Cross-contamination between the environment and domestic animals then progresses through mass transportation and slaughter<sup>68</sup>. Subsequently, wild birds including pigeons and sparrows, rodents, cats, dogs, and insects may be contaminated by contact or ingestion of spilled meat meal, feather meal and other animal by-products outside rendering departments at slaughtering plants, poultry houses and open trucks. These may lead


to contamination of effluents, surface waters, creeks, lakes, rivers, pastures, and soils to colonization of birds, cattle, pigs, sheep, and horses as well as to contamination of animal feeds, or direct re-colonization of farm animals<sup>11,24,54,67,87</sup>. Reportedly however, contamination of the environment by *S. Enteritidis* from meat waste of domestic sources going into the refuse system may be small compared to animal waste and human sewage<sup>68</sup>. Figure 1 presents contamination routes of *S. Enteritidis* in the hen house environment.

### Recent advances in the search for effective vaccines to *S. Enteritidis*

Following the dramatic increase in the last two decades in prevalence of *S. Enteritidis* worldwide, it is becoming a leading cause of food borne illnesses and serious attempts have been made to examine potency of many candidate vaccines to *S. Enteritidis*. There is evidence that administration of live *Salmonella* bacterium orally to newly hatched chicks results in extensive gut colonization and a strong adaptive immunity often leading to rapid protective effects within 24 hours<sup>94</sup>. In this process, establishment and colonization by other bacteria is inhibited by competitive exclusion<sup>96</sup> probably due to competition for binding sites. The presence of a large number of bacteria originating from a live *Salmonella* vaccine in the intestine can also induce infiltration of polymorphonuclear cells into intestinal walls and confer resistance to invasion and systemic spread by virulent *Salmonella* strains.

A recent report suggested the role of SEF14 fimbrial protein in adhesion of *S. Enteritidis* to the host<sup>60</sup>. *Sefa* gene which encodes the subunit of the SEF14 fimbrial protein was cloned, ligated into a temperature sensitive expression vector and transformed into an avirulent strain of *Escherichia coli*. The recombinant strain was used as a vaccine to elicit specific immune responses against the SefA protein of *S. Enteritidis* in one-day old chickens and was re-isolated from the intestines of treated birds for up to 21 days thereafter. Using ELISA, IgA against SefA protein was detected in intestinal secretions from treated birds at 7 days and in bile samples from 14 to 21 days post treatment. Untreated birds did not show evidence of intestinal colonization by the recombinant strain or anti-SefA IgA response in their bile or intestinal secretions.

Additional studies evaluated the effect of oral *S. Enteritidis* and *S. Typhimurium* vaccine (metabolic drift mutants produced by chemical mutagenesis) on colonization of reproductive tract and internal egg contamination among laying hens<sup>35</sup>. Three groups of 30 laying hens were vaccinated orally at day 1, 6 weeks and 16 weeks of age with one or a combination of both vaccine strains and a fourth group (control) was not vaccinated. Birds were intravenously



Broken arrows denote probable but infrequently reported routes of contamination. Full arrows indicate frequently reported routes.

**Fig. 1. *Salmonella* Enteritidis: contamination routes in the hen-house environment**

challenged at 24 weeks of age with 0.5 mL of inoculum containing  $5 \times 10^7$  cfu of *S. Enteritidis* PT4 S1400/94. At three weeks post challenge, the number of oviducts from which *Salmonella* was isolated was significantly lower in the vaccinated than in the non-vaccinated hens. Twelve of 105 (11.4%) batches of eggs were contaminated in birds vaccinated with either vaccine compared to 28 of 105 (26.7%) batches in the unvaccinated group. Contamination was much lower at only 1 of 105 (1.0%) in hens vaccinated with both vaccine strains. These data support the thesis that live vaccines could be valuable in controlling intestinal colonization and internal egg contamination with *S. Enteritidis*. There is compelling evidence that an iron-restricted *S. Enteritidis* PT4 strain 109 vaccine is effective in laying chicken following intramuscular injections and intravascular challenge<sup>102</sup>. Since *S. Enteritidis* spreads through a flock primarily via faecal contamination, vaccines would be considered effective if it increased specific antibody levels in the digestive tract and reduced the amounts of *S. Enteritidis* shed in feces while preventing egg contamination and overall disease transmission by limiting invasion of the bird's internal organs such as the ovaries where eggs are infected internally.

While killed vaccines stimulate only humoral immunity, live *Salmonella* vaccines have the capacity to stimulate both cell-mediated and humoral immunities. Table 1 presents a

list of recent attempts by different laboratories to evaluate efficacy of *S. Enteritidis* vaccine candidates.

### Limitations to complete eradication of *S. Enteritidis* from the hen house

Although identification of eggs as the main source of *S. Enteritidis* food poisoning outbreaks suggested that effective control could be achieved by intensive sampling of birds and diversion of suspect eggs from the market, farmers, public health officials and retailers have been confounded by the intractable nature of the problem of *S. Enteritidis*<sup>12,14,83</sup>. The *S. Enteritidis* pandemic has been most frustrating because it involves the interaction of the pathogen with multiple environments including the hen house (presence of rodents, insects, wild birds, poor ventilation, and accumulation of dust in the hen house), poultry and eggs, as well as the human host. Thus while many laboratories have put in place elaborate efforts to control *Salmonella* infections through vaccinations, creation of a *S. Enteritidis* free hen house remains evasive. This is confounded by the fact that (i) infected adult chickens may remain asymptomatic while producing *Salmonella* contaminated eggs, (ii) infections can be transmitted through cracks on egg shells (iii) horizontal infection is possible among birds in a flock, and (iv) infections are associated with environmental factors such

**Table 1. Recent attempts at evaluation of Salmonella Enteritidis vaccine candidates**

Source	Vaccine type	Methods	Observation
De Buck et al. 2005 <sup>22</sup>	Purified type 1 fimbriae from <i>S. Enteritidis</i> .	Chickens immunized intra-peritoneally at 18 and 21 weeks of age; Three weeks later, immunized and non-immunized birds ( $n = 18$ ) were challenged intravenously with $2 \times 10^7$ live <i>S. Enteritidis</i> .	IgG and IgA were found in eggs and sera of immunized birds. Higher infections in oviducts and <i>Salmonella</i> contamination in eggs of non immunized birds
Betancor et al. 2005 <sup>9</sup>	AroC derivative (LVR02) of Uruguay strain <i>S. Enteritidis</i> .	Oral administration to newly hatched chicks and second dose at 15 days post-hatching.	Protective immunity to oral challenge with <i>S. Enteritidis</i> . Systemic and intestinal infection prevented. Shedding of challenge strain in birds faeces significantly reduced.
De Buck et al. 2004 <sup>21</sup>	Parent strain <i>S. Enteritidis</i> Or <i>fimD</i> mutant (without type 1 fimbriae).	Chickens inoculated intravenously with <i>fimD</i> mutant or its parent strain.	<i>fimD</i> mutant present in blood 3 weeks after infection but wild-type parent strain was cleared in the first 3 days. Eggs of birds infected with <i>fimD</i> were less frequently contaminated with <i>Salmonella</i> , but shells of eggs were more frequently contaminated with the wild-type strain than with the mutant.
Babu et al. 2004 <sup>6</sup>	Live and killed <i>S. Enteritidis</i> .	Chickens 1 <sup>st</sup> immunized at 2 weeks of age followed by a booster dose at 4 weeks, then challenged with <i>S. Enteritidis</i> 2 weeks later (6 weeks of age) and tested for CMI and antibody responses to <i>S. Enteritidis</i> 1 week post-challenge (7 weeks of age).	<i>S. Enteritidis</i> flagellae and con A induced higher splenic cell proliferation. <i>S. Enteritidis</i> shedding was lower in live vaccine group. Splenic CD3 significantly lower and B cells were higher in control group compared to <i>S. Enteritidis</i> challenged group. Serum antibodies to <i>S. Enteritidis</i> flagella and envelope were significantly higher in killed vaccine group.
Khan et al. 2003 <sup>50</sup>	<i>S. Enteritidis</i> outer membrane proteins (OMPs) : 75.6 and 82.3kDa proteins	Twelve 9-wk-old specific-pathogen-free chickens were put in 3 groups of four (I&II immunized subcutaneously with 10 µg of 82.3 kDa, 75.6 kDa, respectively and III; not immunized). Immunized chickens were boosted twice with the same amount of proteins at two week intervals and challenged one week after last boost with 1 ml of $8 \times 10^8$ cfu <i>S. Enteritidis</i> culture. All chickens were sacrificed 48 h after the challenge.	Immunization with OMPs 75.6 and 82.3 kDa proteins led to reduced colonization of <i>S. Enteritidis</i> in intestinal mucosa of chicken. Sera from immunized birds reacted with the 76.5 and 82.3 kDa proteins and not unimmunized sera
Fukutome et al. 2001 <sup>32</sup>	Liposome-associated ultrasonicated whole cell extracts of <i>S. Enteritidis</i> PT4 strain 582	Eight-week-old chickens were immunized with liposome-associated or liposome-free <i>S. Enteritidis</i> cell lysates either intraocularly or intranasally (200 µg or 4 mg protein/100 µL into both eyes or nasal cavities or the same doses in 4 mL orally into the gizzard using a vinyl catheter). Immunization was repeated twice or thrice at 2-week intervals and peripheral blood and intestinal mucosa were collected 1 or 2 weeks after the last immunization.	Liposome-associated antigen induced Serum IgA, IgG, and IgM, but immunization with antigen alone induced only IgG in the intestines. Higher responses were obtained with intra-ocular immunization followed by intra-nasal then oral immunization.
Cooper et al. 1994 <sup>16</sup>	Wild-type <i>S. Enteritidis</i> , LA5 and a genetically defined <i>S. Enteritidis</i> <i>aroA</i> vaccine candidate CVL30 (attenuated in BALB/c mice).	Newly hatched chicks were orally dosed with $10^9$ cfu of bacteria.	Wild-type <i>S. Enteritidis</i> LA5 caused death of 1 in 25 chicks, and gross pathologic symptoms, including pericarditis and perihepatitis in 6 of the 24 survivors after an oral dose of $10^9$ cfu. <i>S. Enteritidis</i> <i>aroA</i> attenuated in BALB/c mice was not virulent.

cfu: colony-forming units; MI: cell-mediated immunity.

as presence of wild birds, insects, rodents in or around the hen house, and farm material. These factors are further confounded by globalization of food supplies, inadvertent introduction of pathogens into new geographic areas by shipping eggs or chickens, movement of travelers, refugees and immigrants, and changing lifestyles with more people eating out at restaurants, canteens, fast food outlets, and street vendors. Insects such as cockroaches and house-flies, as well as chicken droppings can easily get into drinking and feeding trays. Also, wild birds using the hen house as shelter, contamination with human sewage and other wastes, poor ventilation, and high dust levels are believed to aid dissemination of bacteria among chickens by colonization of mucosal surfaces i.e. nares, and conjunctiva<sup>46</sup>. Of the rodents, the house mouse, *Mus musculus* is a rich source of *S. Enteritidis*<sup>20,38,40,42</sup> and the possibility that it may enrich both chicken and human pathogenic strains in the hen house is real.

Although bacteriological screening of hen houses can help to identify flocks at risk<sup>19,42,44</sup>, it is equally difficult to isolate *S. Enteritidis* from hen houses even if the flock is producing contaminated eggs<sup>14,19,33,41,77</sup>. Previous reports have suggested that *S. Enteritidis* infection can be decreased in poultry using competitive exclusion by inoculating hatching chicks with beneficial bacterial cultures<sup>65,70</sup>. Also, moulting, a temporary cessation in egg production induced in older birds by withholding feed and water<sup>23</sup> has been suggested as a strategy that can alter commensal gut flora of poultry. Although this practice increases productivity and extends useful life of the flock, it leads to increased shedding of *Salmonella* in feces<sup>45,46</sup> and risk analysis has suggested that it may double the incidence of egg contamination<sup>91</sup>. According to Kogut et al. (1999)<sup>55</sup>, moulting is stressful to birds.

### Limitations to faster development of viable vaccines

Despite the elaborate efforts by many laboratories to produce viable vaccines to *Salmonella* infection (Table 1), serious limitations remain to be addressed. The fact that chicks are essentially gnotobiotic when hatched and have gastrointestinal tracts filled with meconium makes them particularly susceptible to colonization of the gastrointestinal tract by *Salmonellae* and other microflora<sup>73</sup> and may affect attempts to control colonization by live oral vaccines<sup>16</sup>.

Although unrelated studies have suggested that inactivated *Salmonella* vaccines can protect experimental animals against salmonellosis<sup>28,56,58</sup>, there are contradictions in the literature relating to their efficiency. These contradictions partly arise from variations in formulations, adjuvants,

methods of bacterial inactivation, and animal models used thereby making it difficult to compare results<sup>8</sup>. That inactivated vaccines would lead to serum antibody responses but fail to elicit cell mediated immune responses which are considered important for long term protection and secretory IgA which is needed for protection of mucosal surfaces raises more questions about their efficacy. Use of live vaccines on the other hand raises fears that residual virulence could enter the human food chain or that the public may not readily accept use of genetically manipulated organisms in the food producing animals. Also, given the relatively longer time required to develop and register new live vaccines before use and the threat of rapid emergence and decline of different serovars as seen in *S. Enteritidis* PT4, there is a possibility that the *Salmonella* problem for which a vaccine is developed may no longer exist by the time the vaccine is licensed for use in the field<sup>8</sup>. The risk that residual virulence may also result in vertical transmission and affect productivity cannot be overstated. On the contrary, use of inactivated or subunit vaccines overcome these problems in that they are relatively quick to produce, are stable, and do not contaminate the environment with genetically modified micro-organisms. While fully defined genetic deletions are preferable in live vaccines for purposes of quality control, attenuated pathogens though attractive lack the virulence of live pathogens<sup>25</sup>. Also, differences in types of antigens used, dosing, routes of administration, timing, as well as age and genotypes of chicken used<sup>6,32,35,60</sup> (Table 1) have slowed consensus on the way forward. In addition, while it is not clear whether data obtained from environmentally naive day-old or young chicks can be applicable to chickens of all ages, data obtained from *in vitro* experiments in complete exclusion of *in vivo* regulatory mechanisms may arguably not be entirely representative of the *in vivo* situation.

### Suggested methods for reducing *S. Enteritidis* infection and transmission

Although the data so far generated are of good scientific merit, the threat of *S. Enteritidis* remains and consensus on data from different laboratories using different approaches (Table 1) is hard to achieve. That *S. Enteritidis* was found in sheds after cleaning and disinfection<sup>21</sup> suggests that total elimination from poultry houses would be a daunting task. However, infection rates can be reduced substantially if multi-sectional approaches are embraced by researchers, environmental law enforcement agencies, public health departments, breeders, food handlers including transporters, those working at slaughter houses, retailers, restaurant workers, and consumers. The following approaches could be considered.

- (i) Establishment of inter-laboratory research groups

each addressing specific questions. These studies should be consistent with regard to genotypes of chickens used, study environments, age, sex, feed composition, vaccine types and quantities administered per unit weight of chicks, and routes and timing of vaccine delivery, as well as the types of data collected. Regular peer review of research findings at appropriate symposia would facilitate consensus on management strategies. This approach would be effective especially in areas where chickens or chicks are traded.

(ii) Since vaccinations do not always provide a 100% guarantee that a disease would be prevented from causing losses and widespread use of antibiotics may lead to resistance, breeders should be trained on cleaning and disinfection, use of drag swabs to monitor *Salmonella* contamination before re-stocking poultry houses and hatcheries, regular cleaning of ventilation inlets and fans, occupational health inspections and cool storage to limit growth of *Salmonella* in eggs, and observance of environmental hygiene and prompt vaccination. Testing of feeds especially those of animal origin, and testing to identify positive flocks should be conducted regularly.

(iii) Strict enforcement of environmental safety laws governing transportation of livestock to slaughter houses, hygienic handling at processing plants, safe transportation of offal meats to consumer outlets, prompt sterilization of sewage, dirt removal, and safe disposal of wastes should be observed.

(iv) Because *S. Enteritidis* passes through the food chain from primary production to food service establishments, institutions, and house-holds, there is a need to educate food handlers including those working at slaughter houses, processing factories, transporters, and retailers on good manufacturing practices. Home-based consumers should be cautioned to cook poultry products adequately before consumption.

(v) Constructing a prompt reporting system of new *S. Enteritidis* isolates for researchers would counter emerging threats and facilitate epidemiological analysis. Actually, PulseNet of the Centers for Disease Control and Prevention in the USA (<http://www.cdc.gov/pulsenet/>) and PulseNet Europe (<http://www.pulsenet-europe.org/>) provide useful information on *S. Enteritidis*. A similar approach is under consideration in Japan<sup>98</sup>. These databases provide standardized molecular subtyping (or “fingerprinting”) of food borne disease-causing bacteria by pulsed-field gel electrophoresis.

## Conclusion

Because poultry products are regularly consumed by humans, the risks of *S. Enteritidis*-derived salmonellosis with the attendant public health and economic problems are real. Proper management practices can lead to significant

reduction in the degree of transmission and infections. Inter-laboratory approaches to vaccine development and training of breeders on environmental management at and around hen houses can lead to significant reductions in transmission and overall infections. Good manufacturing practices, hygienic handling of poultry products at processing plants and retail outlets as well as strict adherence to environmental protection laws governing safe transportation of livestock to slaughter houses and offal meat to consumer outlets should be observed. Prompt and efficient sewage treatment and safe waste disposal, as well as hygienic handling and adequate cooking of poultry products before consumption can also limit spread of infections.

## References

1. Altekruze, S. et al. (1993) A comparison of *Salmonella enteritidis* phage types from egg-associated outbreaks and implicated laying flocks. *Epidemiol. Infect.*, **110**, 17-22.
2. Anderson, E. S. & Wilson, E. M. J. (1961) Die Bedeutung der *Salmonella typhimurium*-phagen-typisierung in der Human und Veterinarmedizin. *Zentralblatt für Bakteriologie, Microbiologie und Hygiene series I. Abteilung Orig.*, **224**, 368-373.
3. Anderson, E. S. et al. (1977) Bacteriophage-typing designations of *Salmonella typhimurium*. *J Hygiene, Cambridge*, **78**, 297-300.
4. Anon. (1995) *Salmonella enteritidis* pilot project progress report. A cooperative effort of the Pennsylvania Poultry Producers, Pennsylvania Poultry Federation, Egg Association of America, Pennsylvania Department of Agriculture, Pennsylvania State University, and the United States Department of Agriculture, Washington DC, 48-49.
5. Anon. (2000) Public meeting on *Salmonella enteritidis* research. In *Transcripts of proceedings*, September 8, Hapesville, GA, Food and Drug Administration, USDA, Washington, DC.
6. Babu, U. et al. (2004) *Salmonella enteritidis* clearance and immune responses in chickens following *Salmonella* vaccination and challenge. *Vet. Immunol. Immunopathol.*, **101**, 251-257.
7. Bakersville, A. et al. (1992) Airborne infection of laying hens with *Salmonella enteritidis* phage type 4. *Vet. Rec.*, **130**, 395-398.
8. Barrow, P. A. & Wallis, T. S. (2003) Vaccination against *Salmonella* infections in food animals: rationale, theoretical basis and practical application. In *Salmonella in domestic animals*, eds. Wray, C. & Wray, A., CABI Publ., 323-339.
9. Betancor, L. et al. (2005) An attenuated *Salmonella enteritidis* strain derivative of the main genotype circulating in Uruguay is an effective vaccine for chickens. *Vet. Microbiol.*, **107**, 81-89.
10. Board, R. G. (1996) The course of microbial infection of the hen's eggs. *J. Applied Bacteriol.*, **29**, 319-341.
11. Borland, E. D. (1975) *Salmonella* infection in poultry. *Vet. Rec.*, **97**, 406-408.
12. Byrd, J. A. et al. (1999) Evaluation of *Salmonella* serotype distributions from commercial broiler hatcheries and grower

- houses. *Avian Dis.*, **43**, 39-47.
13. Bryan, E. D. & Doyle, M. P. (1995) Health risks and consequences of *Salmonella* and campylobacter jejuni in raw meat. *J. Food Prot.*, **58**, 326-344.
  14. Caldwell, D. J. et al. (1995) Evaluation of persistence and distribution of *Salmonella* serotype isolation from poultry farms using drag-swab sampling. *Avian Dis.*, **39**, 617- 621.
  15. Callow, B. R. (1959) A new phage typing scheme for *Salmonella typhimurium*. *J. Hygiene, Camb.*, **57**, 346-559.
  16. Cooper, G. L. et al. (1994) Invasiveness and persistence of *Salmonella enteritidis*, *Salmonella typhimurium* and genetically defined *S. enteritidis* AroA strain in young chickens. *Infect. Immunol.*, **62**, 4739-4746.
  17. Davies, R. H. & Wray, C. (1995) Mice as carriers of *Salmonella enteritidis* on persistently infected poultry units. *Vet. Record*, **137**, 337-341.
  18. Davies, R. H. & Wray, C. (1996a) Determination of an effective sampling regime to detect *Salmonella enteritidis* in the environment of poultry units. *Vet. Microbiol.*, **50**, 117-127.
  19. Davies, R. H. & Wray, C. (1996b) Persistence of *Salmonella enteritidis* in poultry units and poultry food. *Brit. Poultry Sci.*, **37**, 589-596.
  20. Davies, R. H. & Wray, C. (1996c) Studies of contamination of three broiler breeder houses with *Salmonella enteritidis* before and after cleansing and disinfection. *Avian Dis.*, **40**, 626-633.
  21. De Buck, J. et al. (2004) Effect of type 1 fimbriae of *Salmonella enterica* serotype Enteritidis on bacteraemia and reproductive tract infection in laying hens. *Avian Pathol.*, **33**, 314-320.
  22. De Buck, J. et al. (2005) Protection of laying hens against *Salmonella enteritidis* by immunization with type 1 fimbriae. *Vet. Microbiol.*, **105**, 93-101.
  23. Deitch, E. A., Winterton, J. & Berg, R. (1987) Effect of starvation on malnutrition and trauma on the gastrointestinal tract flora and bacterial translocation. *Arch. Surg.*, **122**, 1019-1024.
  24. Diesch, S. L. (1978) Environmental aspects of salmonellosis. In Proceedings of national salmonellosis seminar, United States Animal Health Association and Co-sponsors, Washington DC, pp.11.
  25. Dougan, G. et al. (1993) The genetics of *Salmonella* and vaccine development. In *Biology of salmonella*, eds. Cabello, F. et al., Plenum Press, New York, 323-332.
  26. Ebel, E. D., David, M. J. & Mason, J. (1992) Occurrence of *Salmonella enteritidis* in the U.S. commercial egg industry: report on a national spent hen survey. *Avian Dis.*, **36**, 646-54.
  27. Edwards, P. R. (1958) Salmonellosis: observations and control. *Annals of the New York Acad. Sci.*, **70**, 598-613.
  28. Eisenstein, T. K., Killar, L. M. & Sultz, B. M. (1984) Immunity to infection with *Salmonella typhimurium*: mouse-strain differences in vaccine and serum-mediated protection. *J. Infect. Dis.*, **150**, 425-435
  29. Felix, A. & Callow, B. R. (1943) Typing of paratyphoid B bacilli by means of Vi bacteriophage. *Brit. Med. J.*, **ii**, 127-130.
  30. Fisher, I. (2001) Enter-net quarterly *Salmonella* report. ([www2.phls.co.uk/reports/latest.html](http://www2.phls.co.uk/reports/latest.html)). Public Health Laboratory Service.
  31. Forsythe, R. H., Ross, W. J. & Ayres, J. C. (1967) *Salmonella* recovery following gastrointestinal and ovarian inoculation in the domestic fowl. *Poult. Sci.*, **46**, 849-855.
  32. Fukutome, K. et al. (2001) Intestinal mucosal immune response in chickens following intraocular immunization with liposome-associated *Salmonella enterica* serovar *enteritidis* antigen. *Dev. Comp. Immunol.*, **25**, 475-484.
  33. Fuzihara, T. O., Fernandes, S. A. & Franco, B. D. (2000) Prevalence and dissemination of *Salmonella* serotypes along the slaughtering process in Brazilian small poultry slaughter houses. *J. Food Prot.*, **63**, 1749-1753.
  34. Galton, M. M., Steele, J. H. & Newell, K. W. (1964) Epidemiology of salmonellosis in the United States. In *The world of salmonellosis*, ed. van Oye, E., W. Junk Publishers, The Hague, 421-444.
  35. Gantois, I. et al (2006) Oral immunization of laying hens with live vaccine strains of TAD *Salmonella vac E* and TAD *Salmonella vac T* reduces internal egg contamination with *Salmonella enteritidis*. *Vaccine*, **24**, 37-39.
  36. Gauger, H. C. & Greaves, R. E. (1946) Isolation of *Salmonella typhimurium* from drinking water in an infected environment. *Poult. Sci.*, **25**, 476-478.
  37. Gast, R. K. & Holt, P. S. (2000) Deposition of phage type 4 and 13a *Salmonella enteritidis* strains in the yolk and albumen of eggs laid by experimentally infected hens. *Avian Dis.*, **44**, 706-710.
  38. Guard-Petter, J. et al. (1997) On-farm monitoring of Mouse-Invasive *Salmonella enterica* serovar Enteritidis and a model for its association with the production of contaminated eggs. *Applied and Environmental Microbiol.*, **63**, 1588-1593.
  39. Guard-Petter, J. (2001) The chicken, the egg and *Salmonella enteritidis*. *Environmental Microbiol.*, **3**, 421- 430.
  40. Henzler, D. J. & Opitz, H. M. (1992) The role of mice in the epizootology of *Salmonella enteritidis* infection on chicken layer farms. *Avian Dis.*, **36**, 625- 631.
  41. Henzler, D. J. et al. (1994) *Salmonella enteritidis* in eggs from commercial chicken layer flocks implicated in human outbreaks. *Avian Dis.*, **38**, 37-43.
  42. Henzler, D. J., Kradel, D. C. & Sischo, W. M. (1998) Management and environmental risk factors for *Salmonella enteritidis* contamination of eggs. *Am. J. Vet. Res.*, **59**, 824-829.
  43. Higgins, R. et al. (1982) Studies on the dissemination of *Salmonella* in broiler-chicken flocks. *Avian Dis.*, **26**, 26-33.
  44. Hogue, A. et al. (1997) Epidemiology and control of egg associated *Salmonella enteritidis* in the United States of America. *Rev. Sci. Tech.*, **16**, 542- 553.
  45. Holt, P. S. et al. (1994) Effect of two different molting procedures on *Salmonella enteritidis* infection. *Poult. Sci.*, **73**, 1267-1275.
  46. Holt, P. S., Mitchell, B. W. & Gast, R. K. (1998) Airborne horizontal transmission of *Salmonella enteritidis* in molted laying chickens. *Avian Dis.*, **42**, 45-52.
  47. Humphrey, T. J. et al. (1992) Infection of laying hens with *Salmonella enteritidis* PT4 by conjunctival challenge. *Vet. Rec.*, **131**, 386-388.
  48. Humphrey, T. J. (1994) Contamination of egg shell and contents with *Salmonella enteritidis*: a review. *Int. J. Food Microbiol.*, **21**, 31-40.
  49. Keller, L. H. et al. (1995) *Salmonella enteritidis* colonization of the reproductive tract and forming and freshly laid eggs of

- chickens. *Infection and Immunity*, **63**, 2443-2449.
50. Khan, M. I., Fadl, A. A. & Venkitanarayanan, K. S. (2003) Reducing colonization of *Salmonella enteritidis* in chicken by targeting outer membrane proteins. *J. Applied Microbiol.*, **95**, 142-145.
  51. Kinde, H. et al. (2000) Pathologic and bacteriologic findings in 27-week-old commercial laying hens experimentally infected with *Salmonella enteritidis*, phage type 4. *Avian Dis.*, **44**, 239-248.
  52. Kinde, H. et al. (1996a) *Salmonella enteritidis*, phage type 4 infection in a commercial layer flock in Southern California: bacteriologic and epidemiological findings. *Avian Dis.*, **40**, 665-671.
  53. Kinde, H. et al. (1996b) Sewage effluent: likely source of *Salmonella enteritidis*, phage type 4 infections in a commercial chicken layer flock in Southern California. *Avian Dis.*, **40**, 672-676.
  54. Kirkwood, J. K. et al. (1994) *Salmonella enteritidis* excretion by carnivorous animals fed on day-old chicks. *Vet. Rec.*, **134**, 683.
  55. Kogut MH, Genovese KJ & Stanker LH (1999) Effect of induced molting on heterophil function in White Leghorn hens. *Avian Dis.*, **43**, 538-548.
  56. Kuusi, N. et al. (1981) Immunization with outer membrane protein (porin) preparations in experimental murine salmonellosis: effect of lipopolysaccharide. *Infection and Immunity*, **34**, 328-332.
  57. Lamont, S. J., Kaiser, M. G. & Liu, W. (2002) Candidate genes for resistance to *Salmonella enteritidis* colonization in chickens as detected in a novel genetic cross. *Vet. Immunol. Immunopathol.*, **87**, 423-428.
  58. Lindberg, A. A. et al. (1974) Effect of synthetic disaccharide-protein conjugate as immunogen in *Salmonella* infection in mice. *Infection and Immunity*, **10**, 541-545.
  59. Liu, W., Kaiser, M. G. & Lamont, S. J. (2003) Natural resistance-associated macrophage protein 1 gene polymorphisms and response to vaccine against or challenge with *Salmonella enteritidis* in young chicks. *Poult. Sci.*, **82**, 259-266.
  60. Lopes, V. C. et al. (2006) Preliminary evaluation of the use of the *sefA* fimbrial gene to elicit immune response against *Salmonella enterica* serotype *enteritidis* in chickens. *Avian Dis.*, **50**, 85-190.
  61. McDermid, A. S. & Lever, M. S. (1996) Survival of *Salmonella enteritidis* PT4 and *Salmonella typhimurium* Swindon in aerosols. *Letters in Applied Microbiology*, **23**, 107-109.
  62. McIlroy, S. G. & McCracken, R. M. (1990) The current status of the *Salmonella enteritidis* control programme in the United Kingdom. In Proceedings of the 94<sup>th</sup> annual meeting of the United States Animal Health Association, Carter Printing, Richmond, Virginia, 450-462.
  63. Mead, P. S. et al. (1999) Food-related illness and death in United States. *Emerg. Infect. Dis.*, **5**, 607-625.
  64. Methner, U., al-Shabibi, S. & Meyer, H. (1995) Experimental oral infection of specific pathogen-free laying hens and cocks with *Salmonella enteritidis* strains. *Zentralbl. Veterinarmed. [B]*, **42**, 459-469.
  65. Methner, U. et al. (1997) Comparative study of the protective effect against *Salmonella* colonisation in newly hatched SPF chickens using live attenuated *Salmonella* vaccine strains, wild-type *Salmonella* strains or a competitive exclusion product. *Int. J. Food Microbiol.*, **35**, 223-230.
  66. Morris, G. K. et al. (1969) A study of the dissemination of salmonellosis in a commercial broiler chicken operation. *Am. J. Vet. Res.*, **30**, 1413-1421.
  67. Morse, E. V. (1978) Salmonellosis and pet animals. In Proceedings of the national seminars, United States Animal Health Association and other Co-sponsors, Washington, DC, pp.6.
  68. Murray, C. J. (2003) Environmental aspects of *Salmonella*. In *Salmonella* in domestic animals, eds. Wray, C. & Wray, A., CABI Publ., 265-283.
  69. Nakamura, M. et al. (1994) Horizontal transmission of *Salmonella enteritidis* and effect of stress on shedding in laying hens. *Avian Dis.*, **38**, 282-288.
  70. Nuotio, L. et al. (1992) Use of competitive exclusion to protect newly-hatched chicks against intestinal invasion and colonization by *Salmonella enteritidis* PT4. *Br. Poult. Sci.*, **33**, 775-779.
  71. Oosterom, J. (1991) Epidemiological studies and proposed preventive measures in the fight against human salmonellosis. *Int. J. Food Microbiol.*, **12**, 41-52.
  72. PHLS (1989) First report on *Salmonella* in eggs. PHLS Evidence, London: Agricultural Committee of the House of Commons, UK, 14-53.
  73. Popiel, I. & Turnbull, P. C. B. (1985) Passage of *Salmonella enteritidis* and *Salmonella thompson* through chick ileocecal mucosa. *Infect. Immunol.*, **47**, 786-792.
  74. Poppe, C. et al. (1991) The prevalence of *Salmonella enteritidis* and other *Salmonella* spp. among Canadian registered commercial chicken broiler flocks. *Epidemiol. Infect.*, **107**, 201-211.
  75. Poppe, C. (2003) Salmonellosis infections in the domestic fowl. In *Salmonella* in domestic animals, eds. Wray, C. & Wray, A., CABI Publ., 107-132.
  76. Rabsch, W., Tschape, H. & Baumler, A. J. (2001) Nontyphoidal salmonellosis: emerging problems. *Microbes Infect.*, **3**, 237-247.
  77. Riemann, H. et al. (1998) A survey of *Salmonella* by drag swabbing manure piles in California ranches. *Avian Dis.*, **42**, 67-71.
  78. Rigby, C. E. & Pettit, J. R. (1979) Some factors affecting *Salmonella typhimurium* infection and shedding chickens raised on litter. *Avian Dis.*, **23**, 442-455.
  79. Schaaf, J. (1936) Die salmonellose (infektiöse Enteritis, Paratyphose) des Gelflügels ihre Bedeutung und Bekämpfung. *Zeitschrift für Infektionskrankheiten, Parasitäre Krankheiten und Hygiene der Haustiere*, **49**, 322-332.
  80. Shivaprasad, H. L. (2000) Fowl typhoid and pullorium disease. *Rev. Sci. Tech.*, **19**, 405-424.
  81. Snoeyenbos, G. H. et al. (1969) Dynamics of *Salmonella* infection in chicks reared on litter. *Avian Dis.*, **13**, 72-83.
  82. Snoeyenbos, G. H. (1991) Pullorium disease. In *Disease of poultry*, ed. Calnek, B. W., Iowa State University Press, Ames, 73-86.
  83. Soerjadi-Liem, A. S. & Cumming, R. B. (1984) Studies on the incidence of *Salmonella* carriers in broiler flocks entering a poultry processing plant in Australia. *Poult. Sci.*, **63**, 892-895.
  84. Sojka, W. J. et al. (1975) Incidence of *Salmonella* infection in animals in England & Wales, 1968-73. *Vet. Rec.*, **96**, 280-

- 287.
85. St Louis, M. E. et al. (1988) The emergence of grade A eggs as a major source of *Salmonella enteritidis* infections. New implications for the control of salmonellosis. *JAMA*, **259**, 2103-2107.
  86. Stokes, J. L., Osborne, W. W. & Bayne, H. G. (1956) Penetration and growth of *Salmonella* in egg shells. *Food Res.*, **21**, 510-518.
  87. Tannock, G. W. & Smith, J. M. B. (1971) Studies on the survival of *Salmonella typhimurium* and *Salmonella bovis-morbificans* on pasture and in water. *Australian Vet. J.*, **47**, 557-559.
  88. Telzak, E. E. et al. (1990) Anosocomial outbreak of *Salmonella enteritidis* infection due to the consumption of raw eggs. *New Engl. J. Med.*, **32**, 394-397.
  89. Timoney, J. F. et al. (1989) Egg transmission after infection of hens with *Salmonella enteritidis* phage type 4. *Vet. Rec.*, **125**, 600-601.
  90. Ullmann, R. & Scholtze, H. K. (1989) *Salmonella* occurrence in the Erfurt district. *Z Gesamte Hyg.*, **35**, 676-679.
  91. USDA (1989) *Salmonella enteritidis* risk assessment: egg shells and egg products. US Dept. of Agriculture, Washington, DC.
  92. Van de Giessen, A. W. et al. (1992) The identification of *Salmonella enteritidis*-infected poultry flocks associated with an outbreak of human salmonellosis. *Epidemiol. Infect.*, **109**, 405-411.
  93. Van Hemert, S. et al. (2006) Gene expression responses to *Salmonella* infection in the chicken intestine differ between lines. *Vet. Immunol. Immunopathol.*, **114**, 247-258.
  94. Van Immerseel, F. et al. (2002) The effect of vaccination with a *Salmonella* Enteritidis *aroA* mutant on early cellular responses in caecal lamina propria of newly-hatched chickens. *Vaccine*, **20**, 3034-3041
  95. Van Immerseel, F. et al. (2004) Cats as a risk for transmission of antimicrobial drug resistant *Salmonella*. *Emerg. Infect. Dis.*, **10**, 2169-2174.
  96. Van Immerseel, F. et al. (2005) Vaccination and early protection against non-host-specific *Salmonella* serotypes in poultry: exploitation of innate immunity and microbial activity. *Epidemiol. Infect.*, **133**, 959-978.
  97. Ward, L. R., De Sa, J. D. H. & Rowe, B. (1987) A phage-typing scheme for *Salmonella enteritidis*. *Epidemiol. Infect.*, **99**, 291-294.
  98. Watanabe, H. et al. (2002) PulseNet Japan: surveillance system for the early detection of diffuse outbreak based on the molecular epidemiological method. *Kansenshogaku Zasshi*, **76**, 842-848 [In Japanese].
  99. WHO (2007) Antigenic formulae of the *Salmonella* serovars 2007, 9<sup>th</sup> ed. [http://www.pasteur.fr/sante/clre/cadrecnr/salmoms/WKLM\\_2007.pdf](http://www.pasteur.fr/sante/clre/cadrecnr/salmoms/WKLM_2007.pdf).
  100. WHO (2005) Fact sheet No. 139 (Revised April, 2005). <http://www.who.int/mediacentre/factsheets/fs139/en/print.html>.
  101. Williams, J. E., Dillard, L. H. & Hall, G. O. (1968) The penetration patterns of *Salmonella typhimurium* through the outer structures of chicken eggs. *Avian Dis.*, **12**, 445-446.
  102. Woodward, M. J. et al. (2002) The efficacy of Aalenvac, a *Salmonella enterica*, subsp. Enteritidis iron-restricted bacterine vaccine, in laying chickens. *Avian Pathol.*, **31**, 383-392.