

Propagation and distribution system of healthy seedcane as control measures against sugarcane white leaf disease

Production

Implementation

Item: Sugarcane

Chemical pesticide reduction

Outline

We developed a system and manual on field management techniques for propagation of healthy seedcane to control sugarcane white leaf disease (SCWLD) (Fig. 1), distribution methods for the production, procedures to detect pathogens using the loop-mediated isothermal amplification (LAMP) method, and procedures to produce disease-free seedlings using the growth point culture method.

Background/effect/note

SCWLD is one of the most devastating insect-borne diseases affecting sugarcane production in Asia. We considered that the use of healthy seedcane is highly effective in controlling SCWLD. Hence, a manual for the propagation and distribution of healthy seedcane was developed for sugar mills and institutions that produce and distribute seedcane to farmers (Fig. 2). Verification test demonstrated sufficiently low rate of healthy seedcane even though the number of diseased plants increased up to 10-fold in the third generation (Fig. 3). The insecticides that can be used to control the vector are based on information available in Thailand. Users are advised to check and confirm current pesticide treatment regulations in their respective countries.



Fig. 1. A field abandoned due to widespread sugarcane white leaf disease (SCWLD)




Fig. 2. Overview of the healthy seedcane propagation system




Fig. 3. Results of the healthy seedcane propagation verification test
Observations for two generations showed lower disease prevalence (0-20%, mean 5.8%, median 5%) as compared to newly planted fields.



Technical details:

https://www.jircas.go.jp/en/publication/research_results/2020_b10

Contact

greenasia-ml@jircas.go.jp

Japan International Research
Center for Agricultural Sciences

