

TARC Report

The Scolytidae (Coleoptera) of the Philippines*

BY AKIRA NOBUCHI

Forest Protection Division, Forestry and Forest Products Research Institute
(Kukizaki, Inashiki, Ibaraki, 305 Japan)

The Scolytidae, bark beetles and ambrosia beetles, contains a great number of species which are more destructive to forest trees and newly felled timbers than any other group of insect pests. The bark beetles are usually secondary insect pests but, occasionally under conditions favorable for rapid breeding some species such as *Ips calligraphus* in Luzon may increase to large numbers so that they attack healthy trees and become important primary insect pests. On the other hand, the ambrosia beetles or pin-hole borers cause not only mechanical damage to timber, but also provide avenues for the invasion of blue stain or wood rotting fungi from the entrance holes.

As far as the author knows, the fauna of the Philippine Scolytidae contains 277 species belonging to 36 genera¹⁻¹⁰). It is not too much to say that the survey of its fauna is incomplete in comparison with that of Japanese fauna. In this paper, a key to genera and a check list of Scolytidae from the Philippines are presented.

A part of this work was done when the author stayed in the College of Forestry, University of the Philippines at Los Baños as a visiting scientist sent by the Tropical Agri-

culture Research Center in 1978. Thanks are due to the Dean of College and his staffs for kind assistance extended to this study.

Key to genera of Philippine Scolytidae**

1. Pronotum with lateral margins emarginated in basal half; in female pronotum with a conspicuous elongate pore before center; maxillary lobes with both spines and setae on inner edge; xylomycetophagous (Scolytoplatypinae)
.....*Scolytoplatypus* Schaufuss
- Pronotum with lateral margins not emarginated in basal half, without central pore in both sexes; maxillary lobes with either spines or setae on inner edge 2
2. Abdominal sternites strongly ascending; phloepagous 3
- Abdominal sternites not strongly ascending; usually horizontal 4
3. Antennal funicles 4-segmented; antennal clubs solid*Scolytogenes* Eichhoff
- Antennal funicles 6-segmented; antennal clubs with a sutures
.....*Scolytomimus* Blandford
4. Head suboblong and prominent, partially visible from above; pronotum nearly evenly punctured, not asperate on anterior portion; basal margin of elytra usually elevated and transversely crenulate (except Hylastini) (Hylesininae) . . 5
- Head globose, not visible from above, almost entirely concealed by pronotum;

* This article constitutes the "Studies on the Scolytidae XXV" in a series of the author's papers.

** *Coriacephilus* Schedl in Cryphalini and *Ozopemon* Hagedorn in Cryptugini are omitted in this key, because the original descriptions are not enough to include them in this key.

- pronotum usually distinctly roughened, with backwardly directed asperities on anterior portion (except a few species of *Poecilips* in Crypturgini); basal margin of elytra smooth, neither elevated nor crenulate (*Ipinae*) 12
5. Fore tibiae strongly projected outwardly on outer apical angle 6
- Fore tibiae with a row of several tooth-like processes on outside, not projected on outer apical angle 7
6. Eyes divided into two parts; antennal funicles 7-segmented; antennal clubs multi-segmented, appearance by two sutures and some rows of setae; phloeophagous *Sphaerotrypes* Blandford
- Eyes elongate, undivided into two parts; antennal funicles 6-segmented; antennal clubs solid; phloeophagous *Cladoctonus* Strohmeyer
7. Fore coxae rather widely separated at base by a broad prosternal process 8
- Fore coxae almost touching each other *Cladoctonus* Strohmeyer.
8. Maxillary lobes with spine-like setae on inside; antennal clubs with septum or annulate; phloeophagous 9
- Maxillary lobes with hair-like setae on inside; antennal clubs solid; antennal funicles 7-segmented; xylophagous *Dactylipalpus* Chapuis
9. Eyes distinctly emarginated in anterior margin, the emargination extending to middle of eyes; pronotum devoid of asperities; antennal clubs with two septa; antennal funicles 5-segmented *Phloeosinus* Chapuis
- Eyes not emarginated in anterior margin; pronotum usually with a few asperities on antero-lateral portion; antennal funicles 5- or 7-segmented 10
10. Antennal funicles with an oblique septum, 5-segmented *Phloeoditica* Schedl
- Antennal funicles without septum .. 11
11. Antennal funicles 5-segmented; elytra curved down abruptly before apex; abdominal sternites horizontal *Xylechinus* Chapuis
- Antennal funicles 7-segmented; elytra gradually descending behind middle; abdominal sternites weakly ascending behind *Hylesinus* Fabricius
12. Pronotum with comparatively few, large, isolated asperities, with a distinctly elevated summit behind center; elytra usually covered with fine squamiform setae; antennal clubs compressed; phloeophagous, spermatophagous, or xylophagous (*Cryphalini*) 13
- Pronotum not or covered with numerous, small asperities; usually with a weakly elevated summit (except *Dryocoetiops* in Crypturgini and a few species of *Xyleborus* in Xyleborini with a rather strongly elevated summit); elytra usually devoid of squamiform setae (except a few species of *Xyleborus* in Xyleborini) 21
13. Antennal clubs with a distinct septum 14
- Antennal clubs devoid of septum ... 15
14. Antennal clubs with an oblique septum on one side, no suture indicated by row of setae, evenly rounded on sides; fore tibiae rather broad, dilated in middle .. *Cryphalomorphus* Schaufuss
- Antennal clubs with narrow septum between 1st and 2nd sutures, three distinct sutures indicated by rows of setae, sinuate on sides; fore tibiae slender, gradually increasing in width distally *Hypothenemus* Westwood
15. Antennal clubs with sutures indicated by rows of setae 16
- Antennal clubs devoid of suture 19
16. Tarsi with 3rd segment stout or emarginated; antennal funicles 4-segmented 17
- Tarsi with 3rd segment simple; antennal funicles 3-segmented *Margadillius* Hopkins
17. Antennal funicles 4-segmented 18
- Antennal funicles 5-segmented *Hypocryphalus* Hopkins
18. Antennal funicles with 4th segment distinctly wider than the 3rd, nearly as wide as the 2nd; antennal clubs with

- three recurved sutures on anterior face, three procurved sutures on posterior face *Cryphalus* Erichson
- Antennal funicles with 4 segment nearly as wide as the 3rd, narrower than the 2nd; antennal clubs with four sutures, the 4th obscure *Piperius* Hopkins
19. Antennal funicles 3-segmented *Erioschidias* Schedl
- Antennal funicles 4-segmented 20
20. Elytral striae and strial punctures distinct; in male 5th abdominal sternite devoid of tooth *Ptiopodius* Hopkins
- Elytral striae and strial punctures obscure; in male 5th abdominal sternite with four rather prominent teeth *Stephanorhopalus* Hopkins
21. Metepisternum largely covered by elytra, only anterior portion visible; antennal clubs compressed; pronotum finely marginated in basal margin; phloeophagous, xylophagous. (Pityophthorini) *Pityophthorus* Eichhoff
- Metepisternum distinctly visible for entire length; pronotum usually not emarginated in basal margin (except *Coccotrypes* in Crypturgini); if basal margin emarginated, antennal clubs subtruncate 22
22. Maxillary lobes with radiating spines on inner edge; middle and hind tibiae rather slender, abruptly narrowing apically, covered with some teeth outside; anterior plate of proventriculus well developed. Male similar to female in size and general shape (except *Poecilips* and *Coccotrypes* in Crypturgini); phloeophagous, spermatophagous 23
- Maxillary lobes pilose, without radiating spine or inner edge; middle and hind tibiae rather broadly dilated in middle, gradually narrowing apically, armed with many teeth; anterior plate of proventriculus very small and nearly vestigial. Male differs from female in size and general shape; spermatophagous, xylo-mycetophagous 30
23. Elytral declivity with teeth on lateral sides; pronotum with short transverse asperities arranged in concentric and parallel rows on anterior half, usually punctured or granulate on posterior third (Ipinii) 24
- Elytral declivity devoid of tooth; pronotum punctured or irregularly covered with asperities; phloeophagous, spermatophagous, xylophagous (Crypturgini) 25
24. Antennal clubs compressed with strongly procurved sutures ornamented by long setae, basal one of which reaches apical third *Acanthotomicus* Blandford
- Antennal clubs obliquely truncate or flattened, with straight or bisinuate sutures, basal one of which does not reach middle *Ips* Degeer
25. Basal suture of antennal clubs strongly procurved, the shining area semicircular *Ozodendron* Schedl
- Basal suture of antennal clubs weakly recurved or straight, the basal shining area winecup-shaped 26
26. Pronotal summit distinctly elevated; pronotum distinctly declivous in front, rather sparsely intermixed with wide and narrow asperities *Dryocoetiops* Schedl
- Pronotal summit absent or indistinct; pronotum not or slightly declivous in front, with fine and dense asperation, the asperities devoid in some species of *Poecilips* 27
27. Pronotum marginated in posterior margin, serrate in anterior margin *Coccotrypes* Eichhoff
- Pronotum not marginated in posterior margin, not serrate in anterior margin 28
28. Antennal funicles 4-segmented *Cyrtogenius* Strohmeier
- Antennal funicles 5-segmented 29
29. Pronotum acute at least basal half of lateral margins *Poecilips* Schaufuss
- Pronotum not acute in lateral margins *Dryocoetes* Eichhoff
30. Hind tibiae and tarsi strongly compressed and broadly dilated (Eccoapterini)

-*Eccoptopterus* Motschulsky
 - Hind tibiae not strongly compressed or broadly dilated; hind tarsi not compressed 31
 31. Fore tibiae strongly narrowing toward apex (Webbini) *Webbia* Hopkins
 - Fore tibiae broader toward apex (*Xyleborini*) 32
 32. Antennal clubs with two sutures on posterior face; abdominal sternies with sides nearly straight and acuminate from middle toward apex
*Coptoborus* Hopkins
 - Antennal clubs without suture on posterior face; abdominal sternites with sides rounded from middle to apex .. 33
 33. Antennal funicles 4-segmented
*Cnestus* Sampson
 - Antennal funicles 5-segmented 34
 34. Fore tibiae with ventral sides rugose ..
*Arixyleborus* Hopkins
 - Fore tibiae with ventral sides smooth ..
 35
 35. Fore coxae widely separated from each other at base*Xylosandrus* Reitter
 - Fore coxae approximately contiguous at base*Xyleborus* Eichhoff

A check list of Philippine Scolytidae

- Dactylipalpus niger* Schedl: *Philip. J. Sci.*, 90, 87 (1961).
Dactylipalpus transversus Chapuis: *Synopsis des Scolytides*, 220 (1869).
Dactylopalpus unctus Wood: *The Great Basin Naturalist* 21, 8 (1961).
Sphaerotrypes bangensis Eggers: *Philip. J. Sci.*, 33, 73 (1927).
Sphaerotrypes biseriatus Schedl: *Reichenbachia* 4, 243 (1964).
Sphaerotrypes boettcheri Eggers: *Philip. J. Sci.*, 33, 72 (1927).
Sphaerotrypes insularis Eggers: *Philip. J. Sci.*, 33, 74 (1927).
Sphaerotrypes inermis Browne: *Kontyû, Tokyo*, 49, 127 (1981).
Sphaerotrypes moseri Eggers: *Philip. J. Sci.*, 33, 73 (1927).
Sphaerotrypes palawanus Eggers: *Philip. J. Sci.*, 33, 74 (1927).
Sphaerotrypes philippinensis Strohmeier: *Philip. J. Sci.*, 6, 18 (1911).
Diamerus batoensis Eggers: *Treubia*, 9, 392 (1927).
Diamerus curvifer (Walker): *Ann. Mag. nat. Hist.*, (3) 3, 261 (1859).
Diamerus granulatus Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 131 (1923).
Diamerus griseopubescens Schedl: *Tijd. v. Ent.* 93, 52 (1950).
Diamerus interstitialis (Lea): *Proc. r. Soc. Vict. N.S.*, 22, 145 (1910).
Diamerus merinjaki Sampson: *Ann. Mag. nat. Hist.*, (9) 4, 107 (1919).
Diamerus puncticollis Eggers: *Philip. J. Sci.*, 33, 68 (1927).
Diamerus striatus Eggers: *Philip. J. Sci.*, 33, 67 (1927).
Cladoctonus banosus (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 141 (1923).
Phloeoditica phloeosinoides Browne: *Ent. Medd.*, 34, 243 (1966).
Xylechinus formosanus Schedl: *Philip. J. Sci.*, 57, 479 (1935).
Hylesinus birmanus (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 138 (1923).
Hylesinus despectus Walker: *Ann. Mag. nat. Hist.*, (3), 3, 261 (1859).
Hylesinus nigrinus Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 133 (1923).
Hylesinus porcatus Chapuis: *Synopsis des Scolytides* (1869).
Hylesinus sumatranus Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 134 (1923).
Phloeosinus asper (Sampson): *Ann. Mag. nat. Hist.*, (9) 7, 35 (1927).
Phloeosinus australis nagaensis Schedl: *Tijd. v. Ent.*, 82, 30 (1939).
Phloeosinus imitans Eggers: *Philip. J. Sci.*, 33, 75 (1927).
Phloeosinus nonseptis Schedl: *Kol.-forstl. Mitt.*, 5, 173 (1942).
Scolytogenes major (Eggers): *Philip. J. Sci.*, 33, 69 (1927).
Scolytogenes minor (Eggers): *Philip. J. Sci.*, 33, 69 (1927).
Scolytomimus philippinensis (Eggers): *Zool.*

- Meded. R. Mus. nat. Hist. Leiden*, 7, 143 (1923).
- Scolytomimus pusillus* (Eggers): *Philip. J. Sci.*, 33, 88 (1927).
- Erioschidias minimus* (Eggers): *Philip. J. Sci.*, 33, 75 (1927).
- Erioschidias philippinensis* Schedl: *Kontyú*, 35, 126 (1967).
- Margadillius confusus* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 38 (1915).
- Margadillius erythrinae* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 38 (1915).
- Margadillius margadilaonis* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 38 (1915).
- Margadillius minutus* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 37 (1915).
- Ptilopodius minutissimus* Schedl: *Entomol. Blätter*, 39, 34 (1943).
- Ptilopodius rugosus* Schedl: *Entomol. Blätter*, 39, 34 (1943).
- Ptilopodius stephegynis* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 11 (1915).
- Stephanorhopalus melodorum* Schedl: *Entom. Abh. Mus. Tierk. Dresden*, 35, 19 (1966).
- Piperius pini* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 39 (1915).
- Cryphalus boettcheri* Schedl: *Tijd. v. Ent.*, 93, 57 (1950).
- Cryphalus brevisetosus* Schedl: *Entomol. Blätter*, 39, 36 (1943).
- Cryphalus capucinus* Schedl: *Philip. J. Sci.*, 67, 425 (1938).
- Cryphalus densepilosus* Schedl: *Entomol. Blätter*, 39, 36 (1943).
- Cryphalus dilutus* Eichhoff: *Ratio, descriptio, emendatio eorum Tomicinorum*, 490 (1879).
- Cryphalus hagedorni* Eggers: *Entomol. Blätter*, 4, 217 (1908).
- Cryphalus indicus* Eichhoff: *Stettin. ent. Ztg.*, 39, 384 (1878).
- Cryphalus longipilus* Schedl: *Entomol. Blätter*, 39, 34 (1943).
- Cryphalus malloti* Schedl: *Entomol. Blätter*, 39, 37 (1943).
- Cryphalus negrosensis* Browne: *Philip. J. Sci.*, 106, 85 (1977).
- Cryphalus nigricans* Schedl: *Entomol. Blätter*, 39, 35 (1943).
- Cryphalus palawanus* Schedl: *Kol.-forstl. Mitt.*, 5, 174 (1942).
- Cryphalus pusillus* Schedl: *Entomol. Blätter*, 39, 38 (1943).
- Cryphalus resiniferi* Schedl: *Entomol. Blätter*, 39, 36 (1943).
- Cryphalus squamulosus* Strohmeier: *Philip. J. Sci.*, 6, 20 (1911).
- Cryphalus subgranulatus* Schedl: *Entomol. Blätter*, 39, 37 (1943).
- Cryphalus walkeri* Blandford: *Trans. ent. Soc. London*, 200 (1896).
- Cryphalomorphus ater* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 142 (1923).
- Cryphalomorphus bangensis* Eggers: *Philip. J. Sci.*, 33, 75 (1927).
- Cryphalomorphus minor* Eggers: *Philip. J. Sci.*, 33, 76, (1927).
- Cryphalomorphus parvus* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 11 (1915).
- Cryphalomorphus pityophthorinus* (Schedl): *Entomol. Blätter*, 39, 39 (1943).
- Hypothenemus birmanus* (Eichhoff): *Berl. ent. Z.*, 39, 384 (1871).
- Hypothenemus diptercarpi* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 17 (1915).
- Hypothenemus eruditus* Westwood: *Trans. ent. Soc., London*, 1, 34 (1833).
- Hypothenemus glabripennis* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 32 (1915).
- Hypothenemus hispidus* (Eggers): *Sbornik entom. odd. Nár. Musea v Praze*, 3, 156 (1925).
- Hypothenemus psidii* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 32 (1915).
- Hypothenemus pygmaeus* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 24 (1915).
- Hypothenemus setosus* (Eichhoff): *Die forst- und baumzuchtsschädlichen Borkenkäfer*, 81 (1867).
- Hypothenemus sterculiae* (Hopkins): *U.S. Dept., Agr., Off. Sea.*, 99, 32 (1915).
- Hypothenemus tamarindi* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 27 (1915).
- Hypothenemus webbi* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 17 (1915).
- Hypocryphalus bakeri* (Eggers): *Philip. J. Sci.*, 33, 77 (1927).

- Hypocryphalus bidentatus* Browne: *Kontyû, Tokyo*, 49, 383 (1981).
- Hypocryphalus brevior* Schedl: *Entomol. Blätter*, 39, 40 (1943).
- Hypocryphalus kalambanganus* Schedl: *Entomol. Blätter*, 39, 39 (1943).
- Hypocryphalus mindoroensis* Schedl: *Entomol. Blätter*, 39, 39 (1943).
- Hypocryphalus minor* Schedl: *Entomol. Blätter*, 39, 40 (1943).
- Hypocryphalus obesus* (Hopkins): *U.S. Dept. Agr., Off. Sea*, 99, 42 (1915).
- Hypocryphalus obscurus* Hopkins: *U.S. Dept. Agr., Off. Sea*, 99, 42 (1915).
- Hypocryphalus perminimus* (Schedl): *Tijd. v. Ent.*, 85, 13 (1942).
- Hypocryphalus rotundus* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 41 (1915).
- Hypocryphalus striatus* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 42 (1915).
- Coriacephalus cribripennis* Schedl: *Entomol. Blätter*, 39, 40 (1943).
- Cyrtogenius brevior* (Eggers): *Philip. J. Sci.*, 33, 86 (1927).
- Cyrtogenius elongatus* (Eggers): *Philip. J. Sci.*, 33, 85 (1927).
- Cyrtogenius minor* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 218 (1923).
- Cyrtogenius nitidus* (Hagedorn): *Dtsch. ent. Z.*, 1 (1910).
- Cyrtogenius rugicollis* (Eggers): *Tijd. v. Ent.*, 83, 132 (1940).
- Dryocoetiops laevis* (Strohmeyer): *Philip. J. Sci.*, 6, 22 (1911).
- Coccotrypes carpophagus* (Hornung): *Stettin. ent. Ztg.*, 3, 116 (1842).
- Poecilips acuminatus* Schedl: *Entom. Abh. Mus. Tierk. Dresden*, 35, 34 (1966).
- Poecilips advena* (Blandford): *Trans. ent. Soc. London*, 100 (1894).
- Poecilips depressus* (Eggers): *Philip. J. Sci.*, 33, 82 (1927).
- Poecilips incognitus* Schedl: *Philip. J. Sci.*, 90, 89 (1961).
- Poecilips fallax* Eggers: *Treubia*, 9, 399 (1927).
- Poecilips graniceps* (Eichhoff): *Dtsch. ent. Z.*, 21, 120 (1877).
- Poecilips longior* Eggers: *Philip. J. Sci.*, 33, 83 (1927).
- Poecilips medius* Eggers: *Philip. J. Sci.*, 33, 84 (1927).
- Poecilips myristicae* (Roepke): *Treubia*, 1, 24 (1919).
- Poecilips oblongus* Eggers: *Philip. J. Sci.*, 33, 83 (1927).
- Poecilips papuanus* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 148 (1923).
- Poecilips philippiensis* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 145 (1923).
- Poecilips striatus* (Eggers): *Philip. J. Sci.*, 33, 82 (1927).
- Poecilips subcristosus* (Blandford): *Trans. ent. Soc. London*, 224 (1896).
- Poecilips vulgaris* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 151 (1923).
- Ozopemon angustae* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 159 (1923).
- Ozopemon aplanatus* Schedl: *Kol.-forstl. Mitt.*, 5, 177 (1942).
- Ozopemon brownei* Schedl: *Ann. Mag. nat. Hist.*, (12) 6, 297 (1953).
- Ozopemon diperocarpi* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 49 (1915).
- Ozopemon latus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 156 (1923).
- Ozopemon laevis* Strohmeyer: *Philip. J. Sci.*, 6, 22 (1911).
- Ozopemon major* Strohmeyer: *Philip. J. Sci.*, 6, 23 (1911).
- Ozopemon obanus* Hagedorn: *Dtsch. ent. Z.*, 3 (1910).
- Ozopemon parinari* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 48 (1915).
- Dryocoetes lusonicus* Schedl: *Entomol. Blätter*, 39, 41 (1943).
- Dryocoetes schultzei* Schedl: *Philip. J. Sci.*, 90, 88 (1961).
- Ozodendron papuanus* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 157 (1923).
- Arixyleborus confinus* (Eggers): *Philip. J. Sci.*, 33, 106 (1927).
- Arixyleborus dipterocarpi* Browne: *Kontyû, Tokyo*, 49, 133 (1981).
- Arixyleborus granulifer* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 206

- (1923).
- Arixyleborus hirsutulus* Schedl: *Kontyû*, 37, 212 (1970).
- Arixyleborus imitator* (Eggers): *Philip. J. Sci.*, 33, 105 (1927).
- Arixyleborus pusillus* (Eggers): *Philip. J. Sci.*, 33, 108 (1927).
- Arixyleborus rugosipes* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 59 (1915).
- Arixyleborus sublaevis* (Eggers): *Philip. J. Sci.*, 33, 104 (1927).
- Arixyleborus subsimilis* Schedl: *Kontyû*, 38, 362 (1970).
- Coptodryas confusa* Hopkins: *U.S. Dept. Agr., Off. Sea.*, 99, 54 (1915).
- Xyleborus adossuarinus* Schedl: *Philip. J. Sci.*, 80, 367 (1951).
- Xyleborus agnatus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 197 (1923).
- Xyleborus agraphus* Schedl: *Orient. Ins.*, 11, 503 (1977).
- Xyleborus alsapanicus* Schedl: *Tijd. v. Ent.*, 93, 59 (1950).
- Xyleborus amphicranoides latecavatus* Eggers: *Philip. J. Sci.*, 33, 95 (1927).
- Xyleborus amplexicauda* Hagedorn: *Dtsch. ent. Z.*, 9 (1910).
- Xyleborus armillatus* Schedl: *Philip. J. Sci.*, 52, 199 (1933).
- Xyleborus artheybridus* Schedl: *Tijd. v. Ent.*, 93, 66 (1950).
- Xyleborus balbalanus* Eggers: *Philip. J. Sci.*, 33, 95 (1927).
- Xyleborus barbatus* Hagedorn: *Dtsch. ent. Z.*, 11 (1910).
- Xyleborus benguetensis* Schedl: *Tijd. v. Ent.*, 93, 71 (1950).
- Xyleborus bicolor* Blandford: *Trans. ent. Soc. London*, 113 (1894).
- Xyleborus bidentatus* (Motschulsky): *Bull. Soc. Imp. Nat. de Moscou*, 36, 514 (1863).
- Xyleborus cognatus* Blandford: *Ann. Soc. ent. Fr.*, 65, 19 (1896).
- Xyleborus concisus* Blandford: *Trans. ent. Soc. London*, 107 (1894).
- Xyleborus costatomorphus* Schedl: *Ann. Mag. nat. Hist.*, (12) 3, 897 (1950).
- Xyleborus criticus* Schedl: *Ann. Mag. nat. Hist.*, (12) 3, 899 (1950).
- Xyleborus cuneolus* Eggers: *Philip. J. Sci.*, 33, 92 (1927).
- Xyleborus cylindricus* Eggers: *Philip. J. Sci.*, 33, 94 (1927).
- Xyleborus cylindromorphus* Eggers: *Philip. J. Sci.*, 33, 96 (1927).
- Xyleborus desectus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 167 (1923).
- Xyleborus destruens* Blandford: *Trans. ent. Soc. London*, 221 (1896).
- Xyleborus dipterocarpi* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 58 (1915).
- Xyleborus diversicolor* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 202 (1923).
- Xyleborus dossuarinus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 187 (1923).
- Xyleborus eggersianus* (Schedl): *Philip. J. Sci.*, 60, 60 (1936).
- Xyleborus emarginatus* Eichhoff: *Stettin. ent. Ztg.*, 39, 392 (1878).
- Xyleborus erinaceus* Eggers: *Philip. J. Sci.*, 33, 103 (1927).
- Xyleborus exiguus* (Walker): *Ann. Mag. nat. Hist.*, (3) 3, 260 (1859).
- Xyleborus eximus* Schedl: *Kontyû*, 38, 362 (1970).
- Xyleborus exsculptus* Eggers: *Philip. J. Sci.*, 33, 101 (1927).
- Xyleborus fallax* Eichhoff: *Stettin. ent. Ztg.*, 39, 392 (1878).
- Xyleborus fastigatus* Schedl: *Philip. J. Sci.*, 56, 402 (1935).
- Xyleborus ferrugineus* Fabricius: *Syst. Eleuth.*, 2, 388 (1801).
- Xyleborus formicatus* Eichhoff: *Berl. ent. Z.*, 12, 151 (1868).
- Xyleborus funereus* Lea: *Proc. r. Soc. Vict. (N.S.)* 22, 139 (1910).
- Xyleborus funereus robustus* Schedl: *Philip. J. Sci.*, 51, 103 (1933).
- Xyleborus granularis* Schedl: *Ann. Mag. nat. Hist.*, (12) 3, 898 (1950).
- Xyleborus haddeni* Schedl: *Philip. J. Sci.*, 51, 101 (1933).
- Xyleborus hybridus* Eggers: *Philip. J. Sci.*, 33, 90 (1927).
- Xyleborus impar* Eggers: *Philip. J. Sci.*, 33, 89 (1927).
- Xyleborus indicus* Eichhoff: *Stettin. ent.*

- Ztg.*, 39, 392 (1878).
- Xyleborus interjectus* Blandford: *Trans. ent. Soc. London*, 576 (1894).
- Xyleborus intermedius* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 201 (1923).
- Xyleborus leavis* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 201 (1923).
- Xyleborus latecornis* Schedl: *Kontyú*, 37, 212 (1969).
- Xyleborus lewisi* Blandford: *Trans. ent. Soc. London*, 104 (1894).
- Xyleborus limbatus* Schedl: *Philip. J. Sci.*, 60, 65 (1936).
- Xyleborus loricatus* Schedl: *Philip. J. Sci.*, 52, 200 (1933).
- Xyleborus lugubris* Eggers: *Philip. J. Sci.*, 33, 98 (1927).
- Xyleborus macopterus* Schedl: *Stylops*, 4, 271 (1935).
- Xyleborus mancus* Blandford: *Trans. ent. Soc. London*, 428 (1898).
- Xyleborus marginicollis* Schedl: *Philip. J. Sci.*, 60, 64 (1936).
- Xyleborus mascarensis* Eichhoff: *Ratio, descriptio, emendatio eorum Tomycinorum*, 372 (1878).
- Xyleborus melas* Eggers: *Philip. J. Sci.*, 33, 93 (1927).
- Xyleborus metacuneolus* Eggers: *Tijd. v. Ent.*, 83, 150 (1940).
- Xyleborus mindanaensis* Eggers: *Philip. J. Sci.*, 33, 93 (1927).
- Xyleborus minimus* Schedl: *Ent. Arb. Mus. Georg Frey*, 6, 305 (1955).
- Xyleborus mucronatus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 191 (1923).
- Xyleborus mutilatus* Blandford: *Trans. ent. Soc. London*, 103 (1894).
- Xyleborus nepocranus* Schedl: *Tijd. v. Ent.*, 82, 45 (1939).
- Xyleborus obliquesectus* Eggers: *Philip. J. Sci.*, 33, 99 (1927).
- Xyleborus obtusicollis* Schedl: *Philip. J. Sci.*, 67, 427 (1938).
- Xyleborus orbus* Browne: *Ent. Medd.*, 34, 249 (1966).
- Xyleborus pedellus* Schedl: *Kontyú*, 37, 213 (1969).
- Xyleborus perbrevis* Schedl: *Tijd. v. Ent.*, 93, 59 (1950).
- Xyleborus perforans* (Wollaston): *Catalogue of the coleopterous insects of Madeira in the collection of the British Museum*, 96 (1857).
- Xyleborus permarginatus* Schedl: *Philip. J. Sci.*, 52, 200 (1933).
- Xyleborus perpilosellus* Schedl: *Philip. J. Sci.*, 56, 402 (1933).
- Xyleborus persimilis* Eggers: *Philip. J. Sci.*, 33, 97 (1927).
- Xyleborus pertortuosus* Schedl: *Kol.-forstl. Mitt.*, 5, 186 (1942).
- Xyleborus pilipunctatus* Browne: *Ent. Medd.*, 34, 251 (1966).
- Xyleborus pilosulus* Eggers: *Philip. J. Sci.*, 33, 100 (1927).
- Xyleborus polyodon* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 196 (1923).
- Xyleborus posticepilosus* Schedl: *Tijd. v. Ent.*, 93, 92 (1950).
- Xyleborus procerior* Schedl: *Mitt. münch. ent. Ges.*, 32, 189 (1942).
- Xyleborus pruinus* Blandford: *Trans. ent. Soc. London*, 214 (1896).
- Xyleborus pseudocylindricus* Eggers: *Treubia*, 9, 402 (1927).
- Xyleborus pullus* Schedl: *Philip. J. Sci.*, 80, 370 (1952).
- Xyleborus pumilus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 209 (1923).
- Xyleborus pusillus* Schedl: *Philip. J. Sci.*, 90, 92 (1961).
- Xyleborus quadraticollis* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 197 (1923).
- Xyleborus recidens* Sampson: *Ann. Mag. nat. Hist.*, (9) 11, 287 (1923).
- Xyleborus rufobrunneus* Eggers: *Philip. J. Sci.*, 33, 101 (1927).
- Xyleborus rufus* Schedl: *Tijd. v. Ent.*, 93, 74 (1950).
- Xyleborus schultzei* Schedl: *Tijd. v. Ent.*, 93, 68 (1950).
- Xyleborus semiopacus* Eichhoff: *Ratio, descriptio, emendatio eorum Tomycinorum*, 334 (1878).
- Xyleborus semipolitus* Schedl: *Tijd. v. Ent.*, 93, 70 (1950).

- Xyleborus seaspinatus* Schedl: *Stylops*, 4 (12), 271 (1935).
- Xyleborus similis* Ferrari: *Die forst- und baumzuchtschädlichen Borkenkäfer*, 23 (1867).
- Xyleborus sordicauda* (Motschulsky): *Bull. Soc. Imp. Nat. de Moscou*, 36, 514 (1863).
- Xyleborus streblicola* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 55 (1915).
- Xyleborus subagnatus* Schedl: *Reichenbachia*, 3, 314 (1964).
- Xyleborus subcostatus* Eichhoff: *Berl. ent. Ztg.*, 12, 281 (1868).
- Xyleborus subdolosus* Schedl: *Tijd. v. Ent.*, 85, 44 (1942).
- Xyleborus submarginatus* Eggers: *Tijd. v. Ent.*, 83, 150 (1940).
- Xyleborus sublongus* Eggers: *Philip. J. Sci.*, 33, 99 (1927).
- Xyleborus subnaevus* Schedl: *Kontyû*, 34, 40 (1966).
- Xyleborus terminaliae* (Hopkins): *U.S. Dept. Agr., Off. Sea.*, 99, 54 (1915).
- Xyleborus torquatus* Eichhoff: *Berl. ent. Z.*, 12, 146 (1868).
- Xyleborus trispinatus* Browne: *Kontyû*, 49, 130 (1981).
- Xyleborus truncatipennis* Schedl: *Philip. J. Sci.*, 90, 92 (1961).
- Xyleborus tuberculosissimus* Eggers: *Tijd. v. Ent.*, 83, 152 (1940).
- Xyleborus turbineus* Sampson: *Ann. Mag. nat. Hist.*, (9) 11, 288 (1923).
- Xyleborus ursa* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 172 (1923).
- Xyleborus ursinus* Hagedorn: *Dtsch. ent. Z.*, 381 (1908).
- Xyleborus ursulus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 173 (1923).
- Xyleborus ursus* Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 173 (1923).
- Xyleborus validus* Eichhoff: *Ann. Soc. ent. Belg.*, 18, 202 (1875).
- Xyleborus vestitus* Schedl: *Ann. Mag. nat. Hist.*, (10) 8, 341 (1931).
- Xyleborus xanthopus* Eichhoff: *Berl. ent. Z.*, 12, 151 (1868).
- Xylosandrus compactus* Eichhoff: *Ann. Soc. ent. Belg.*, 18, 202 (1875).
- Xylosandrus discolor* (Blandford): *Trans. ent. Soc. London*, 429 (1898).
- Xylosandrus luzonicus* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 174 (1923).
- Xylosandrus morigerus* (Blandford): *Insect life*, 6, 260 (1894).
- Xylosandrus pusillus* (Schedl): *Philip. J. Sci.*, 90, 91 (1961).
- Cnestus bicornioides* (Schedl): *Philip. J. Sci.*, 80, 368 (1952).
- Cnestus bimaculatus* (Eggers): *Philip. J. Sci.*, 33, 88 (1927).
- Cnestus nitidus* (Schedl): *Tijd. v. Ent.*, 93, 87 (1950).
- Cnestus punctulatus* (Schedl): *Tijd. v. Ent.*, 93, 61 (1950).
- Webbia dasyurus* Browne: *Kontyû Tokyo*, 49, 133 (1981).
- Webbia dentatus* Eggers: *Philip. J. Sci.*, 33, 108 (1927).
- Webbia dipterocarpi* Hopkins: *U.S. Dept. Agr. Bur., Tech. Ser.*, 17, (2), 223 (1915).
- Webbia mucronatus* Eggers: *Philip. J. Sci.*, 33, 107 (1927).
- Webbia obtusipennis* Schedl: *Kontyû*, 34, 38 (1966).
- Webbia philippinensis* Schedl: *Orient. Ins.*, 5, 374 (1971).
- Webbia platypoides* Eggers: *Philip. J. Sci.*, 33, 105 (1927).
- Webbia 14-spinatus* Sampson: *Ann. Mag. nat. Hist.*, (9) 7, 34 (1921).
- Webbia 18-spinatus* Browne: *Sarawak Mus. J.*, 3, 493 (1958).
- Eccoptopterus collaris* (Eggers): *Zool. Meded. R. Mus. nat. Hist. Leiden*, 7, 194 (1923).
- Eccoptopterus gracilipes* (Eichhoff): *Not. Leyd. Mus.*, 8, 25 (1886).
- Eccoptopterus sagittarius* Schedl: *Tijd. v. Ent.*, 82, 41 (1939).
- Eccoptopterus spinosus* (Olivier): *Entomologie, ou histoire naturelle des insectes*, 3, 9 (1794).
- Acanthotomicus pereziguus* (Blandford): *Trans. ent. Soc. London*, 201 (1896).
- Acanthotomicus tuberculatus* (Eggers): *Philip. J. Sci.*, 33, 79 (1927).

- Ips bicaudatus* Eggers: *Philip. J. Sci.* **33**, 80 (1927).
Ips bispinosus Eggers: *Philip. J. Sci.*, **33**, 78 (1927).
Ips calligraphus (Germar): *Insectorum species novae aut minus cognitae*, 461 (1824).
Pityophthorus philippinensis Schedl: *Philip. J. Sci.*, **60**, 59 (1936).
Scolytoplastypus javanus Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, **7**, 164 (1923).
Scolytoplastypus papuanus Eggers: *Zool. Meded. R. Mus. nat. Hist. Leiden*, **7**, 165 (1923).
Scolytoplastypus pusillus Eggers: *Entomol. Blätter*, **31**, 243 (1935).

References

- 1) Beaver, R. A. & Browne, F. G.: The Scolytidae and Platypodidae (Coleoptera) of Penang, Malaysia, *Orient. Ins.*, **12**, 575-624 (1978).
- 2) Browne, F. G.: Some Scolytidae (Coleoptera) from tropical Africa. *Rev. Zool. Bot. Afr.* **87**, 679-696 (1973).
- 3) Browne, F. G.: Three new species of Scolytidae and Platypodidae (Coleoptera) from Malaysia. *Philip. J. Sci.*, **106**, 85-86 (1977).
- 4) Browne, F. G.: Bark beetles and ambrosia

- beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese Ports, with descriptions of new species, I-V. *Kontyû, Tokyo*, **48**, 370-379, 380-389, 482-489, 490-500 (1980), **49**, 125-136 (1981).
- 5) Nobuchi, A.: Ambrosia beetles found in imported tropical timbers from Southeast Asia and others. *Bull. Gov. For. Exp. Sta.*, **301**, 1-46 (1978) [In Japanese with English summary].
 - 6) Nobuchi, A.: Ambrosia beetles of mahogany in the Philippines, Studies on Scolytidae XV. *Kontyû*, **47**, 406-407 (1979).
 - 7) Nobuchi, A.: Bark and ambrosia beetles of imported timbers, identification of bark beetles found in Siberian and tropical timbers. Ringyo Gijutsu Shinkosho, Tokyo, **79** (1981) [In Japanese].
 - 8) Sato, K.: A list of bark beetles and pin-hole borers imported into Japan with timbers from abroad. *Res. Bull. Pl. Prot. Japan*, Suppl. **12**, 1-67 (1975) [In Japanese with English summary].
 - 9) Schedl, K. E.: Check list of the Scolytidae and Platypodidae from the Philippine Islands. *Entom. Abh. Mus. Tierk. Dresden*, **35**, 1-122 (1966).
 - 10) Schedl, K. E.: Bark and timber beetles of the Oriental region. *Orient. Ins.*, **9**, 451-460 (1975).

(Received for publication, July 14, 1982)