

TARC Report

The Platypodidae (Coleoptera) of the Philippines

By AKIRA NOBUCHI

Forest Protection Division, Forestry and Forest Products Research Institute
(Kukizaki, Ibaraki, 305 Japan)

The Platypodidae consists of more than a thousand species in the world as far as we know, and most species are distributed in the tropical and subtropical region. The fauna of the Philippine Platypodidae is composed of 74 species belonging to eight genera.^{1-4, 6-11} Adults are readily distinguished from the Scolytidae by their larger size, slender and cylindrical bodies; their large exserted heads; rounded and convex eyes; the very long slender tarsus, with the first tarsal segment longer than the second to fifth combined; and the freely articulated sternites.

All members of this family are xylo-myctophagous ambrosia beetles, with a monogamous social organization. The chief damage by these beetles is due to the black-stained galleries (pin-holes) made deeply in the sapwood or heartwood of trees felled for timber. Their damage to green logs are more severe than scolytid ambrosia beetles, because they make usually broader and deeper tunnels into wood. In this paper, I deal a key to genera and a check list of Platypodidae with the literature of their original descriptions from the Philippines. This work is the second part of the study on the Philippine ambrosia beetles as the sequel to the Scolytidae⁵⁾ and was done when I stayed in the College of Forestry, University of Philippines at Los Baños as a visiting scientist sent by the Tropical Agriculture Research Center in 1978. Thanks are due to the Dean of College and his staffs for kind assistance extended to this study.

Key to Philippine subfamilies of Platypodidae

1. Antennal funicles 4-segmented. Pronotum with a median sulcus 2
- Antennal funicles 2-segmented. Pronotum without median sulcus .. Platytarsilinae
2. Fore coxae widely separated from each other at base; outer face of fore tibiae marked by a variable number of transverse carinae in both sexes Diaporinae
- Fore coxae approximated contiguous at base 3
3. Eyes rounded to short oval, strongly convex 4
- Eyes elongate, weakly convex; outer face of fore tibiae marked by a variable number of transverse carinae in both sexes Periommatinae
4. Outer face of fore tibiae marked by a variable number of transverse carinae in male; in female the carina reduced, at least partly replaced by granules.....
- Crossotarsinae
- Outer face of fore tibiae similarly carinæ in both sexes Platypodinae

Key to Philippine genera of subfamily Crossotarsinae

1. Femoral grooves of pronotum angulate at anterior extremity, gently rounded behind *Crossotarsus* Chapuis
- Femoral grooves of pronotum not angulate at anterior extremity, the posterior ex-

tremity angulate or marked by a small tooth *Trachyostus* Schedl

Key to Philippine genera of subfamily Platypodinae

- Elytra of the male usually at least weakly declivous toward apex, rarely longitudinally horizontal and then provided with a vertical apical rim. *Platypus* Herbst
Elytra, at least in male, longitudinally horizontal and not provided with a distinct vertical apical rim *Baiocis* Browne

In subfamily Periommatinae, a single genus *Spathidicerus* Chapuis has been found from the Philippines.

Key to Philippine genera of subfamily Diaporinae

- Antennae inserted in discal and lower part of eyes in both sexes; scutellum usually sunk *Diapus* Chapuis
Antennae inserted in discal and lower part of eyes in male, but inserted in discal upper part of median level of eyes in female; scutellum large, elevated
..... *Genyocerus* Motschulsky

In subfamily Platytarsilinae, a single genus *Platytarsus* Schedl has been found from the Philippines.

A Check List of Philippine Platypodidae

Subfamily Crossotarsinae
Genus *Crossotarsus* Chapuis

Section *Crossotarsi genuini*

- Crossotarsus cincinnatus* Chapuis: *Monographie des Platypides*, 57 (1865).
Crossotarsus comatus Chapuis: *Monographie des Platypides*, 59 (1865).
Crossotarsus laratensis Beeson: *Ind. For. Rec.*, N.S., (3) 3, 56 (1937).
Crossotarsus lecontei Chapuis: *Monographie des Platypides*, 60 (1865).

Crossotarsus mniszechi Chapuis: *Monographie des Platypides*, 62 (1865).

Crossotarsus palatus Beeson: *Ind. For. Rec.*, N.S., (3) 3, 55 (1937).

Crossotarsus piceus Chapuis: *Monographie des Platypides*, 56 (1865).

Section *Crossotarsi ventricorni*

Crossotarsus flavomaculatus Strohmeyer: *Ent. Mitt.*, 1, 40 (1912).

Section *Crossotarsi coleoptrati*

Crossotarsus squamulatus Chapuis: *Monographie des Platypides*, 87 (1865).

Crossotarsus squamulatus fractus Sampson: *Ann. Mag. Nat. Hist.*, (8) 10, 249 (1912).

Crossotarsus squamulatus rastellus Beeson: *Ind. For. Rec.*, N.S., (3) 3: 82 (1937).

Crossotarsus terminatus Chapuis: *Monographie des Platypides*, 85 (1865).

Crossotarsus tribuaricus Schedl: *J. Fed. Mal. St. Mus.*, 18, 358 (1939).

Section *Crossotarsi alternante-depresso*
Crossotarsus octocostatus Schedl: *Philip. J. Sci.*, 56, 398 (1935).

Section *Crossotarsi subdepressi*

Crossotarsus externe-dentatus (Fairmaire): *Rev. et Mag. de Zool.*, 2, 51 (1850).

Crossotarsus saltator Schedl: *Philip. J. Sci.*, 57, 480 (1935).

Section *Crossotarsi incertae sedis*

Crossotarsus concavifrons Schedl: *Philip. J. Sci.*, 56, 399 (1935).

Genus *Trachyostus* Schedl

Trachyostus parvus Browne: *Philip. J. Sci.*, 89, 213 (1960).

Genus *Platypus* Herbst

Section *Platypi hirtelli*
Platypus curtus Chapuis: *Monographie des Platypides*, 261 (1865).
Platypus perrisi Chapuis: *Monographie des Platypides*, 260 (1865).

Section *Platypi antennati*

Platypus excedens Chapuis: *Monographie des Platypides*, 276 (1865).

Platypus nocuus Schedl: *Philip. J. Sci.*, 57, 487 (1935).

Section *Platypi sulcati*

Platypus geminatus Chapuis: *Monographie des Platypides*, 239 (1865).

Platypus jansoni Chapuis: *Monographie des Platypides*, 244 (1865).

Platypus lozonicus Schedl: *Philip. J. Sci.*, 57, 485 (1935).

Platypus setaceus Chapuis: *Monographie des Platypides*, 234 (1865).

Platypus tenellus Schedl: *Philip. J. Sci.*, 57, 485 (1935).

Platypus hybridus Schedl: *Philip. J. Sci.*, 56, 395 (1935).

Platypus spectabilis Schedl: *Philip. J. Sci.*, 56, 396 (1935).

Section *Platypi emarginati*

Platypus deflectus Schedl: *Ent. Ber.*, 10, 355 (1941).

Platypus ptochooides Schedl: *Kontyû*, 38, 367 (1970).

Section *Platypi oxyuri*

Platypus artesolidis Schedl: *Kol.-forstl. Mitt.*, 5, 200 (1942).

Platypus hirtus Schedl: *Ent. Arab., Mus. Frey*, 6, 310 (1955).

Platypus oneratus Schedl: *Kontyû*, 38, 366 (1970).

Platypus solidus Walker: *Ann. Mag. Nat. Hist.*, (3) 2, 286 (1858).

Section *Platypi semidepressi*

Platypus inutilus Schedl: *Philip. J. Sci.*, 57, 479 (1935).

Platypus schultzeanus Schedl: *Ent. Abh. Mus. Tierk. Dresden*, 35, 98 (1967).

Section *Platypini lunati*

Platypus andrewesi (Strohmeyer): *Ent. Bl.*, 6, 128, 131 (1910).

Platypus subdepressus (Schedl): *Philip. J. Sci.*, 56, 397 (1935).

Platypus tayabasi (Schedl): *Philip. J. Sci.*,

57, 481 (1935).

Section *Platypi sulcato-striati*

Platypus philippensis Blandford: *Trans. ent. Soc. London*, 193 (1896).

Section *Platypi signati*

Platypus transformis Schedl: *J. Fed. Mal. St. Mus.*, 18, 16 (1936).

Section *Platypi cupulati*

Platypus agnatus Schedl: *Kontyû*, 37, 217 (1969).

Platypus biuncus Blandford: *Trans. ent. Soc. London*, 194 (1896).

Platypus caliculus Chapuis: *Monographie des Platypides*, 280 (1865).

Platypus cavus Strohmeyer: *Ent. Bl.*, 7/8, 162 (1913).

Platypus cupulatus Chapuis: *Monographie des Platypides*, 278 (1865).

Platypus forficula Chapuis: *Monographie des Platypides*, 283 (1865).

Platypus lepidus Chapuis: *Monographie des Platypides*, 282 (1965).

Platypus noonadanae Browne: *Ent. Medd.*, 34, 237 (1966).

Platypus pallidus Chapuis: *Monographie des Platypides*, 284 (1865).

Platypus pseudocupulatus Schedl: *J. Fed. Mal. St. Mus.*, 17, 635 (1935).

Section *Platypi truncatipenni*

Platypus fallax Schedl: *Kontyû*, 38, 367 (1970).

Section *Platypi pseudospinulosi*

Platypus sexporus (Schedl): *Philip. J. Sci.*, 57, 484 (1935).

Platypus ornaticeps (Schedl): *Philip. J. Sci.*, 56, 401 (1935).

Section *Platypi inaequali-costati*

Platypus shoreanus bifurcus (Schedl): *Philip. J. Sci.*, 67, 422 (1938).

Platypus shoreanus mutilus Schedl: *J. Fed. Mal. St. Mus.*, 18, 359 (1939).

Section *Platypi quadrifissi*

Platypus calignosus Schedl: *Ann. Mus. Civ.*

Genova, 59, 58 (1936).
Platypus quadrifissilis Schedl: *Ann. Mus. Civ. Genova*, 59, 57 (1936).

Section *Platypi adjuncti*

Platypus coleoptratus (Schedl): *Philip. J. Sci.*, 56, 400 (1935).
Platypus loricatus (Sampson): *Ann. Mus. Civ. St. Nat.*, 51, 71 (1923).

Genus *Baiocis* Browne

Biocis anaticeps (Schedl): *Kontyû*, 37, 207 (1969).
Biocis inimicus (Schedl): *Philip. J. Sci.*, 57, 482 (1935).
Biocis pernanulus (Schedl): *Philip. J. Sci.*, 57, 482 (1935).

Subfamily Periommatainae

Genus *Spathidicerus* Chapuis
Spathidicerus philippinensis Schedl: *Kol.-forstl. Mitt.*, 5, 216 (1942).

Subfamily Diaporinae

Genus *Genyocerus* Motschulsky
 (= *Diacarus* Schedl)
Genyocerus abdominalis (Schedl): *Kol.-forstl. Mitt.*, 5, 218 (1942).
Genyocerus biporus (Schedl): *Kol.-forstl. Mitt.*, 5, 217 (1942).
Genyocerus compactus (Schedl): *Kontyû*, 34, 40 (1966).
Genyocerus decemspinatus (Schedl): *Kol.-forstl. Mitt.*, 5, 216 (1942).
Genyocerus multiporus (Schedl): *Philip. J. Sci.*, 60, 67 (1936).
Genyocerus philippinensis (Schedl): *Kol.-forstl. Mitt.*, 5, 216 (1942).
Genyocerus tenellus (Schedl): *Kontyû*, 34, 42 (1966).

Genus *Diapus* Chapuis

Diapus pendleburyi Schedl: *J. Fed. Mal. St. Mus.*, 18, 18 (1936).
Diapus plumatus Schedl: *Philip. J. Sci.*, 60, 66 (1936).
Diapus pusillimus Chapuis: *Monographie des Platypides*, 335 (1865).
Diapus quadrispinosus Schedl: *Kontyû*, 37,

218 (1969).
Diapus quinquespinatus Chapuis: *Monographie des Platypides*, 334 (1865).

Subfamily Platytarsilinae

Genus *Platytarsulus* Schedl

Platytarsulus elongatus Schedl: *Kol.-forstl. Mitt.*, 5, 215 (1942).

References

- 1) Browne, F. G.: Bark Beetles and ambrosia beetles (Coleoptera, Scolytidae and Platypodidae) intercepted at Japanese ports, with descriptions of new species, I, III-VII, *Kontyû*, 48, 370-379, 482-489, 490-500 (1980), 49, 125-136 (1982), 51, 554-572 (1983).
- 2) Nobuchi, A.: Platypodidae found in imported tropical timbers from South Eastern Asia and others. *Bull. Gov. For Exp. Sta.*, 296, 101-155 (1977) [In Japanese with English summary].
- 3) Nobuchi, A.: Ambrosia beetles of mahogany in the Philippines, Studies on Scolytidae XV. *Kontyû*, 47, 406-407 (1979).
- 4) Nobuchi, A.: Bark and ambrosia beetles of imported timbers II, Identification of bark beetles found in Siberian and tropical timbers. *Ringyô Gijutsu Shinkoshô*, Tokyo, 75. (1981) [In Japanese].
- 5) Nobuchi, A.: The Scolytidae (Coleoptera) of the Philippines, *JARQ*, 16, 296-305 (1983).
- 6) Sato, K.: A list of bark beetles and pin-hole borers imported into Japan with timbers from abroad. *Res. Bull. Pl. Prot. Jpn.*, Suppl. 12, 1-67 (1975) [In Japanese with English summary].
- 7) Schedl, K. E.: Check list of the Scolytidae and Platypodidae from the Philippine Islands. *Entom. Abh. Mus. Tierk. Dresden*, 35, 1-122 (1966).
- 8) Schedl, K. E.: Pin-hole borers and bark-beetles intercepted from imported logs in Japanese ports. *Kontyû*, 34, 29-43 (1966).
- 9) Schedl, K. E.: Bark-beetles and pin-hole borers intercepted from imported logs and seeds in Japanese ports II. *Kontyû*, 35, 119-129 (1967).
- 10) Schedl, K. E.: Bark-beetles and pin-hole borers intercepted from imported logs in Japanese ports III-IV. *Kontyû*, 37, 202-219 (1969), 38, 358-370 (1970).
- 11) Schedl, K. E.: Monographie der Familie Platypodidae, Coleoptera. Den Haag, 322 (1972).

(Received for publication, October 17, 1984)