

TARC Notes

Antagonistic iron-deficiency of rice plants growing in acid red latosol in Paraná, Brazil

A distinct chlorosis of rice plants growing in acid red latosol was found in upland fields, just after a long rain, in newly developed lowland fields, or even under greenhouse (pot culture) conditions in Paraná, Brazil. A preliminary test on Fe application to the chlorotic leaves indicated that the symptom is a typical Fe deficiency.

Contents of Fe, Mn, and Zn in the soil, extracted with various extractants, are given in Table 1. Based on data already available^{1,3,4,6,7}, the content of exchangeable Fe extracted with N-ammonium acetate (pH 4.5) was regarded of an adequate level, whereas the contents of exchangeable Mn and easily reducible Mn, extracted with N-ammonium acetate (pH 4.5) and N-ammonium acetate (pH 4.5)+0.2% hydroquinone respectively, were extremely high. Available Zn, extracted with N-ammonium acetate (pH 4.5) or 0.1 N HCl, was also regarded to be at an adequate level.

Since available Mn in soil is influenced by a rise of pH, moisture range, and organic matter content^{2,6}, a factorial pot experiment was carried out in a greenhouse during a

Table 1. Available Fe, Mn and Zn in soil

Extractants	ppm in soil*		
	Fe	Mn	Zn
H ₂ O	0.0	4.3	0.0
N-NH ₄ OAc pH 4.5	14.0	66.8	0.75
N-NH ₄ OAc pH 4.5+ 0.2% hydroquinone	368	500	—
N-KCl pH 7.0	5.0	57.3	1.65
0.1 N-HCl	80.0	90.0	1.55

* Extracted by adding 100 ml of extractant to 10 g of oven-dried soil, and shaking for 1 hr at room temperature. Atomic absorption spectrophotometer of Perkin-Elmer, Model-503, was used for assays.

period from August to October 1976 to know the effects of soil pH and application of Fe and Zn on growth of rice plants in upland as well as lowland conditions. Although the content of available Zn in the soil was found to be sufficient, as shown above, Zn application treatments was included in the experiment, because Fe application might cause an antagonistic depression of Zn uptake. The experiment was run with 3 replications.

IR 305-3-15, one of the varieties which showed severe chlorosis in the field was used. Each pot contained 6 kg of subsurface soil taken from the experimental field of Instituto Agronômico do Paraná with fertilizers (N 1.0g, P₂O₅ 1.0 g and K₂O 0.5 g/pot), and 4 plants were grown per pot. Distilled water was used for watering. Plant growth and contents of Fe, Mn, and Zn in plants were determined at 50 days after germination.

In the upland condition, chlorosis occurred in control plots and Zn plots at pH 4.6 and 5.0 at 15 days after germination. It occurred, though slightly, even in Fe and Fe + Zn plots at every pH level from 29 days after germination. Under the lowland condition, chlorosis appeared in control and Zn plots of every pH level at 12 days after germination, and the symptom became severe as the growth




Fig. 1. Effect of soil pH and micro-nutrient application on rice growth in upland and lowland condition.

LSD (5%): condition=0.21, pH=0.16, micro-nutrient=0.32

Table 2. Effect of soil pH and micro-nutrient application on Zn, Fe and Mn concentration of rice leaves (pot culture experiment)

Treatment of pH	Treatment of micro-nutrients											
	Control			Fe			Zn			Fe+Zn		
	Zn	Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn	Zn	Fe	Mn
ppm in leaves												
Upland condition												
pH 4.6	42	141	6310	34	175	4550	161	125	3580	59	152	3150
5.0	55	143	6140	25	128	4830	148	138	4600	50	163	4640
5.6	29	98	4560	22	120	5380	142	105	5940	46	110	5870
Lowland condition												
pH 4.6	55	152	4100	42	168	4350	161	108	4900	89	158	3890
5.0	52	120	5920	33	137	4800	120	249	4460	48	157	3990
5.6	21	125	3270	21	122	4970	88	147	3480	54	379	3600

Note 1) pH of original soil is 4.6, and amended by CaCO_3 application

2) Fe application: 30 ppm as Fe EDTA

3) Zn application: 5 ppm as ZnSO_4

4) Stems were excluded to avoid contaminations by soil

advanced. It appeared also, though slightly, in Fe + Zn plot of pH 5.6, Fe plot of every pH, and Fe + Zn plot of pH 4.6 and 5.0 at 19, 26, and 35 days after germination respectively.

Fig. 1 shows top growth of plants as effected by the treatments. Variance analysis indicated the significant differences with a extremely larger extent of variance ratios, among the main effects of water condition, pH, and the application of micro-nutrients, although the interactions of water condition \times pH, water condition \times micro-nutrient, and pH \times micro-nutrient were also significant.

Effect of Fe and Fe + Zn applications on plant growth was remarkable at pH 4.6 and 5.0 in both upland and lowland conditions. The combined application of Fe and Zn gave better growth than the application of Fe alone at pH 4.6 and 5.0 in lowland condition and at pH 5.0 in upland condition. However, the application of Zn alone gave no clear effect: it rather caused an inhibitive effect at pH 4.6 in lowland condition.

Contents of Fe, Mn, and Zn in leaves are given in Table 2. Under the upland condition, the application of Fe increased the Fe content, but decreased Mn content (except at pH 5.6) and Zn content. The application of Fe + Zn

caused increased Zn contents in addition to the increased Fe contents and decreased Mn contents. The application of Zn caused remarkable increases in Zn contents with decreased contents of Fe and Mn (except at pH 5.6). The combined application of Fe and Zn resulted in the Zn contents similar to or higher than those of the control plot, and higher Fe contents with lower Mn contents. Under the lowland condition, the similar trend is observed though it is not so clear as above.

The relations observed among Fe, Mn, and Zn contents suggest the antagonistic absorption of them. It can be considered that the increased plant growth in the Fe plot at pH 4.6 and 5.0 is attributable to an increased Fe absorption in relation to Mn absorption. The better growth in the Fe + Zn plot than the Fe plot can be accounted for by the increased absorption of Zn in addition to the increased Fe absorption and decreased Mn absorption. In other words, the increased growth observed in the Fe plot was still limited by the reduced Zn absorption caused by Fe application, so that the Fe + Zn application could give a further increase of growth.

These results strongly suggest that the chlorosis observed in acid red latosol is caused

by an antagonism between Fe and Mn: Fe-deficiency occurs in the presence of an excess of Mn in the soil containing an adequate amount of available iron.

- 1) Adams, F. & Wear, J. I.: Manganese toxicity and soil acidity in relation to crinkle leaf cotton. *Soil. Sci. Soc. Am. Proc.*, 21, 305-308 (1957).
- 2) Christensen, P. D., Toth, S. J. & Bear, F. E.: The status of soil manganese as influenced by moisture, organic matter, and pH. *Soil Sci. Soc. Am. Proc.*, 15, 279-282 (1950).
- 3) Committee of Soil Chemical Analysis, Agr. Forest, and Fisheries Research Council, MAF.: Analytical methods of soil nutrients. pp. 430, Yokendo, Tokyo, Japan (1975) [In Japanese].
- 4) Gangwar, M. S. & Chandra, S. K.: Estimation of critical limit for zinc in rice soils. *Comm. in Soil Sci. and Plant Analysis*, 6, 641-654 (1975).
- 5) Ishizuka, Y. & Ando, T.: Interaction between manganese and zinc in growth of rice plant. *Soil Sci. & Plant Nut.*, 14, 201-206 (1968).
- 6) Sherman, G. D. & Fujimoto, C. K.: The effect of the use of lime, soil fumigants and mulch on the solubility of manganese in Hawaiian soils. *Soil Sci. Soc. Am. Proc.*, 11, 206-210 (1946).
- 7) Wear, J. I. & Evans, C. E.: Relationship of zinc uptake by corn and sorghum to soil zinc measured by three extractants. *Soil Sci. Soc. Am. Proc.*, 32, 543-546 (1968).

Received for publication, February 1, 1978.

Yoshikazu OHNO *Tropical Agriculture Research Center, Japan.*

Celso Jamil MARUR *Instituto Agronômico do Paraná, Brasil.*

Lauro Akio OKUYAMA *Instituto Agronômico do Paraná, Brasil.*