

Infectious Bursal Disease of Chickens

By KATSUYA HIRAI

Faculty of Agriculture, Gifu University

Infectious bursal disease (IBD), formerly termed Gumboro disease had been observed in the U.S.A. towards the end of the 1950 but it was not until 1962 that it was first described by Cosgrove. At first the disease was erroneously called "Avian nephrosis". The disease is a viral disease of young chickens for which the causative virus is tentatively classified as a member of Reoviridae. The virus induces atrophy of the bursa of Fabricius (BF) as a result of necrosis of lymphocytes and also causes a general lymphocidal effect in other lymphoid organs, including the thymus and spleen. The BF has a key role in the development and maturation of the humoral immune response of the chicken. A number of workers have demonstrated that infection with IBD virus at a young age can lead to immunosuppression. Therefore, this disease is of interest immunologically. In the present paper some of our recent studies will be described briefly.

Morphological characterization of the virion^{2,4,7)}

The virus was tentatively classified as a member of Reoviridae. The classification was based upon the morphology under ultrathin-section electron microscopy of the BF from infected chickens and upon limited biochemical characterization. Little information is available on the ultrastructure of the IBD virus. In this section, electron micrographs of purified IBD virus, negatively stained with phosphotungstate, are presented.

IBD virus apparently has an outer layer, as indicated by the large size of intact particles with hexagonal outlines (Plate 1 a-f).

Moreover, the capsomeric detail on the main capsid surface of IBD virus often appeared partially obscured by such a layer. This outer layer was very thin (7 to 8 nm) and continuous, with T-shaped structures suggestive of the configuration reported for rotavirus but not as clearly defined^{12,15,18)}. Although the outer layers of reovirus and orbivirus are also indistinct, they are featureless^{11,18)}. The outer layer of IBD virus may exert some stability to the structure of the underlying major capsid layer. By rotational enhancement of image detail (Plate 2 a-f), the results of this study confirmed our previous report that IBD virus has an icosahedral symmetry of $T=3$, with a probable 32 large capsomeres. This type of symmetry of the unique feature of subunit sharing are morphological characteristics reported for the Reoviridae family^{14,15)}. The results obtained are summarized as follows. An outer layer surrounding the capsid of IBD virus was evident from electron micrographs of intact virus particles having diameters of 62 to 63 nm. The capsid was found to be composed of large morphological units or capsomeres, measuring about 12 nm in diameter. The architecture of the capsid appears to be that of $T=3$ symmetry, with probable 32 morphological units by rotational enhancement of image detail.

Structural proteins of the virion^{3,7)}

Apart from the initial study by Nick et al.¹³⁾, little information is available on the structural proteins of IBD virus, using poly-peptide gel electrophoresis. The exact number of polypeptides and their locations within the virion are not yet clearly understood. The results of this study demonstrated that




Plate 1. Electron micrographs of negatively stained IBD virus particles.

- a. A large number of IBD virus recovered from the 1.34 g/ml CsCl fraction. Bar=100 nm. $\times 45,000$.
- b. Intact virions of IBD virus showing hexagonal outlines. Bar=100 nm. $\times 130,000$.
- c. Intact virions of IBD virus with an outer layer. Bar=50nm. $\times 330,000$.
- d. Higher magnification of an intact virion showing the outer layer. The surface of the particles appear to have T-shaped morphology which is continuous. Bar=50 nm. $\times 450,000$.
- e. IBD virus particles recovered from the 1.35 g/ml CsCl fraction. Intact virions of IBD virus (arrow) and particles with clear capsomeres showing the loss of the outer layer (double arrow). Bar=100 nm. $\times 190,000$.
- f. Particles showing clearly discernible capsomeres. Bar=100 nm. $\times 200,000$.




Plate 2. continued

- a-c. Single IBD virus particle viewed on a 3-fold axis of symmetry representing $n=0$, $n=6$, and $n=5$ rotations, respectively. Enhancement of capsomeres is evident in $n=6$ rotation. Bar = 50 nm. $\times 390,000$.
- d-f. Single IBD virus particle viewed on a 5-fold axis of symmetry representing $n=0$, $n=5$, and $n=6$ rotations, respectively. Enhancement of capsomeres is evident in $n=5$ rotation. Bar = 50 nm. $\times 390,000$.

Table 1. A comparison of molecular weights* of IBD virus proteins with those of rotavirus, bluetongue virus and avian reovirus

IBDV	Human rotavirus	Bluetongue virus	Avian reovirus
133	127	140	140
124	103	110	125
98	97	101	115
	88	82	85
51	58	61	72
33	32	42	40
26.5	26	29	36
23	21		32

* Molecular weight $\times 10^3$

structural proteins of IBD virus consist of seven species, two major and five minor polypeptides. These are P1 to P7, with molecular weights of 133×10^3 , 124×10^3 , 98×10^3 , 51×10^3 , 33×10^3 , 26×10^3 , and 23×10^3 , respectively. The major polypeptides were associated with the smaller subunit particles and string-like structures. Therefore, these polypeptides may be the capsomeral proteins of IBD virus. These results differed from those reported by Nick et al.¹³ with regard to the number of polypeptides and their molecular weights. Only four polypeptides were found in the previous study, three of which did compare closely in molecular weight with our P4, P5, and P6 proteins. However, a polypeptide having a molecular weight of 11×10^3 was not detected in our preparations. The limited amount of purified IBD virus obtained may not have provided sufficient material or detection of all structural polypeptides. As shown in Table 1, the molecular weights of the IBD virus proteins compared more closely with those reported for the human rotavirus. However, the major polypeptide species (P3 and P4) of IBD virus did not correspond with those major proteins of the human rotavirus, nor did IBD virus have an eighth polypeptide of molecular weight 88×10^3 . Our results concerning the similarity of its protein composition to the rotavirus group suggest its tentative inclusion into the Reoviridae family.

Some characterization of immunosuppression in chickens by IBD virus^{1,5,8)}

This disease is of interest immunologically, since the function of the bursa-dependent lymphoid system is affected in young chickens. The mechanism of the immunosuppression is not fully understood, but presumably it results from the loss of immunocompetent lymphocytes. The present studies were undertaken to compare sequential changes in the proportion of B- and T-lymphocytes and measure serum immunoglobulin concentrations in chickens infected with IBD virus at different ages. The results obtained are summarized as follows. Chickens were infected with IBD virus in ovo or at different times posthatching to 6 weeks of age. The B- and T-cell responses in the lymphoid tissues and blood were examined sequentially to 8 weeks of age by using indirect immunofluorescence. The proportion of B-cells was consistently lower in infected birds than in controls, especially in chicks infected at embryos or at 1 day old. The proportion of T-cells increased following these early infections but was slightly lower in spleen and blood of birds infected at 1, 4, and 6 weeks of age. Serum IgM levels dropped significantly after infection, regardless of the time of infection. IgG levels decreased following early infection but increased after infection at 1 week old or more. The results strongly suggest that B-cells are the target for IBD virus infection.

Replication of IBD virus in cultured lymphocytes⁶⁾

Hitchner reviewed studies on laboratory host systems for IBD virus and noted that the virus replicates in chicken embryos and that embryos-adapted virus could be cultured in cell cultures of chicken embryo origin, with consequent cytopathic effects. No published

information is available on replication of virulent, nonadapted strains of the virus in cultured lymphocytes. It has been shown in a previous section that a specific lymphocyte type might serve as the target for IBD virus replication *in vivo*. As part of an investigation of the mechanism by which this virus induces immunosuppression, it was of interest to study virus replication *in vitro*. The results obtained are summarized as follows. The *in vitro* susceptibility of chicken lymphocytes to a wild strain of IBD virus was investigated by using immunofluorescence and virus assays as infection criteria. A variety of Marek's disease lymphoblastoid cell lines, all of thymus (T-cell) origin, were refractory to virus exposure. However, a bursa (B-cell)-derived lymphoblastoid cell line from an avian leukosis virus-induced tumor was highly susceptible. Viral antigen appeared in the cytoplasm of 20 to 30% of the cells, and large amounts of cell-free virus were released, with maximum yields occurring by 3 days postinfection. The virus also replicated in a small percentage of normal lymphocytes prepared from lymphoid tissues and peripheral blood of chickens. Pretreatment of the lymphocytes, with heat-inactivated anti-B-cell serum or with antiserum against fowl immunoglobulin M before inoculating them with the virus blocked the virus infection; no blocking occurred with anti-T-cell serum or with specific antiserum against fowl immunoglobulin G or immunoglobulin A. This suggests that surface immunoglobulin M-bearing B-lymphocytes were the target cells for infection.

Pathogenicity of IBD virus in mouse, rat and hamster^{9,10)}

Rinaldi et al.¹⁷⁾ and Petek et al.¹⁰⁾ reported that egg-adapted strain of IBD virus was pathogenic for suckling mice. Very little is known of the pathogenicity to laboratory animals. The results obtained are summarized as follows. The susceptibility of mice and rats became lower according with the advance

in age regardless of inoculation routes. Mice and rats were the most susceptible to intracerebral inoculation. The infectivity titers in the brains were the highest regardless of the routes of virus inoculation. The titer in the brains increased logarithmically from 12 to 60 hrs postinoculation. A plateau of $10^{6.8}$ to $10^{7.7}$ PFU/0.2 ml was maintained between 60 to 72 hrs. It is also noted that precipitating antigen was produced in the brain of mice and rats inoculated with the virus. The antigen formed 2 or 3 precipitin lines against specific antiserum. The principal changes of the brains were degeneration and destruction of the nerve cells in thalamus and cortex. A specific fluorescence was found in the cytoplasm of nerve cells. The virus particles were observed in the cytoplasm of the infected nerve cells. Suckling hamsters inoculated intracerebrally with the virus developed listlessness and marked weight loss beginning on postinoculation days 12 to 14. The occipital curvature of the skull became prominent at 17 to 19 days after inoculation. All had mild to severe hydrocephalus. The gross hydrocephalus was first seen at day 14 and reached a maximum on the 21 day.

References

- 1) Hirai, K. et al.: Immunodepressive effect on infectious bursal disease virus in chickens. *Avian Dis.*, **18**, 50-57 (1974).
- 2) Hirai, K., Shimakura, S. & Kawamoto, E.: Electron microscope observation of infectious bursal disease virus. *Avian Dis.*, **18**, 467-471 (1974).
- 3) Hirai, K., Kawamoto, E. & Shimakura, S.: Some properties of precipitating antigens associated with infectious bursal disease virus. *Infect. Immun.*, **10**, 1235-1240 (1974).
- 4) Hirai, K. & Shimakura, S.: Structure of infectious bursal disease virus. *J. Virol.*, **14**, 964-974 (1974).
- 5) Hirai, K., Kunihiro, K. & Shimakura, S.: Characterization of the immunosuppression of chickens by infectious bursal disease virus. *Avian Dis.*, **23**, 950-965 (1979).
- 6) Hirai, K. & Calnek, B. W.: In vitro replication of infectious bursal disease virus in established lymphoid cell lines and chicken B lymphocytes. *Infect. Immun.*, **25**, 964-970 (1979).

- 7) Hirai, K., Kato, N. & Shimakura, S.: Further morphological characterization and structural proteins of infectious bursal disease virus. *J. Virol.*, **32**, 323-328 (1979).
- 8) Hirai, K. & Funakoshi, H.: Sequential changes of surface immunoglobulin of B cells in infectious bursal disease virus-infected chickens (Unpublished data).
- 9) Kawamoto, E. et al.: Growth of infectious bursal disease virus in suckling mice and rats. The 81st Meeting of the Jap. Soc. Vet. Sci. (1976) [In Japanese].
- 10) Kawamoto, E. et al.: Hydrocephalus of suckling hamster by infectious bursal disease virus. The 84th Meeting of the Jap. Soc. Vet. Sci. (1977) [In Japanese].
- 11) Luftig, R. B. et al.: An ultrastructural study of virions and cores of reovirus type 3. *Virology* **48**, 171-181 (1972).
- 12) Martin, M. L., Palmer, E. L. & Middleton, P. J.: Ultrastructure of infantile gastroenteritis virus. *Virology*, **68**, 146-153 (1975).
- 13) Nick, H., Cursiefen, D & Becht, H.: Structural and growth characteristics of infectious bursal disease virus. *J. Virol.*, **18**, 227-234 (1976).
- 14) Palmer, E. L. & Martin, M. L.: The fine structure of the capsid of reovirus type 3. *Virology*, **76**, 109-113 (1977).
- 15) Palmer, E. L., Martin, M. L. & Murphy, F. A.: Morphology and stability of infantile gastroenteritis virus: comparison with reovirus and blue-tongue virus. *J. Gen. Virol.*, **35**, 403-414 (1977).
- 16) Petek, M., D'Aprile, N. & Cancellotti, f.: Biological and physico-chemical properties of the infectious bursal disease virus (IBDV). *Avian Path.*, **2**, 135-152 (1973).
- 17) Rinaldi, A. et al.: Untersuchungen über die Ätiologie der sogenannten Gumboro-Krankheit. II. Pathogene Wirkung des Virus auf einige Laboratoriumstiere. 4th Congress of the World Veterinary Poultry Association, Belgrade, 309-313 (1969).
- 18) Stannard, L. M. & Schoub, B. D.: Observations on the morphology of two rotaviruses. *J. Gen. Virol.*, **37**, 435-439 (1977).